



RedNHE

Red Nacional de
Investigadores
en Economía

Rational Bubbles: Too Many to be True?

Martín Solá (Universidad Torcuato di Tella)

DOCUMENTO DE TRABAJO N° 240

Mayo de 2023

Los documentos de trabajo de la RedNIE se difunden con el propósito de generar comentarios y debate, no habiendo estado sujetos a revisión de pares. Las opiniones expresadas en este trabajo son de los autores y no necesariamente representan las opiniones de la RedNIE o su Comisión Directiva.

The RedNIE working papers are disseminated for the purpose of generating comments and debate, and have not been subjected to peer review. The opinions expressed in this paper are exclusively those of the authors and do not necessarily represent the opinions of the RedNIE or its Board of Directors.

Citar como:

Solá, Martín (2023). Rational Bubbles: Too Many to be True?. *Documento de trabajo RedNIE N°240.*

Rational Bubbles: Too Many to be True?

February 1, 2023

Abstract

Issues that arise in the practical implementation of the [Phillips, Wu, and Yu \(2011\)](#) and [Phillips, Shi, and Yu \(2015a\)](#) recursive procedures for identifying and dating explosive episodes in time series are considered. It is argued that the use of critical values for right-tailed unit-root tests obtained under the assumption of a drift whose magnitude depends on the sample size and becomes negligible in large samples results in over-rejection of the unit-root hypothesis when, as in many financial time series, the deterministic drift effect is non-negligible relatively to the stochastic trend. In addition, the standard practice of using conventional levels of significance for critical values involved in the algorithms that locate the origination and termination dates of explosive episodes lead to false discoveries of explosiveness with high probability. The magnitude of these difficulties is quantified via simulations, using artificial data whose properties reflect closely those of real-world time series such as stock prices and dividends. The findings offer a potential explanation for the relatively large number of apparent explosive episodes that are often reported in applied work. Ways of overcoming the aforementioned difficulties by using bootstrap-based calibration techniques are considered. An empirical example focusing on monthly U.S. data on real stock prices and real dividends is also discussed.

Keywords: Bootstrap; Bubbles; Date-stamping; Explosive behavior; Unit-root test.

JEL Classification: C12, C15, C22.

1 Introduction

Empirical detection of rational bubbles has been an active area of research for a long time. This is not very surprising since the possibility of substantial price changes occurring independently of movements in the underlying market fundamentals is of considerable interest from both a theoretical and a policy-making point of view. As a result, various approaches to detecting explosiveness associated with rational bubbles have been proposed in the literature – [Gürkaynak \(2008\)](#) and [Homm and Breitung \(2012\)](#) provide useful overviews and comparisons. The focus in this paper is on the popular and influential procedures introduced in [Phillips, Wu, and Yu \(2011\)](#) (PWY henceforth) and later generalized by [Phillips, Shi, and Yu \(2015a\)](#) (PSY henceforth), which rely on right-tailed unit-root tests based on autoregressive specifications which may exhibit explosive behavior over a subset of the data. The PWY and PSY test statistics are suprema of conventional Augmented Dickey–Fuller (ADF) statistics based on suitably constructed subsamples. These statistics are used both to test the unit-root hypothesis against an explosive alternative and to estimate the beginning and end dates of explosive episodes.¹ The procedures have been applied to a wide variety of time series, including stock prices ([PWY, PSY, Homm and Breitung \(2012\)](#), [Phillips, Shi, and Yu \(2014\)](#), [Phillips and Shi \(2020\)](#), [Monschang and Wilfing \(2021\)](#)), commodity prices ([Phillips and Yu \(2011\)](#), [Gutierrez \(2013\)](#)), [Etienne, Irwin, and Garcia \(2014\)](#), [Fantazzini \(2016\)](#), [Harvey, Leybourne, Sollis, and Taylor \(2016\)](#), [Long, Li, and Li \(2016\)](#)), house prices ([Phillips and Yu \(2011\)](#), [Greenaway-McGrevy and Phillips \(2016\)](#), [Pavlidis, Yusupova, Paya, Peel, Martínez-García, Mack, and Grossman \(2016\)](#), [Hu and Oxley \(2018\)](#)), and cryptocurrency prices ([Cheung, Roca, and Su \(2015\)](#), [Corbet, Lucey, and Yarovaya \(2018\)](#), [Bouri, Shahzad, and Roubaud \(2019\)](#), [Hafner \(2020\)](#)).

The objective of this paper is to highlight and discuss two potential difficulties en-

¹Like many other statistical tests for bubbles, the PWY and PSY procedures are designed to detect bubble-like explosive behavior in a time series. However, it is worth bearing in mind that such behavior may not necessarily be associated with rational bubbles. Economic variables such as asset prices, consumer prices, or foreign-exchange rates are related to underlying fundamental determinants (e.g., dividend payments or money supply). Therefore, any synchronous explosive episodes in the time paths of both prices and the underlying fundamentals would suggest that explosiveness is driven by the fundamentals component rather than by extraneous factors.

countered in the application of the PWY and PSY recursive techniques for testing for the presence of explosive episodes and for identifying their origination and termination dates, both of which can lead to over-detection of explosive behavior. These issues have not attracted the deserved attention either in the theoretical literature, much of which has focused on the properties of the testing and date-stamping procedures in the presence of periodically collapsing bubbles, or in the empirical literature in which the procedures have been applied to real-world data.

The first issue relates to the specification of the data-generating process (DGP) under the null hypothesis being tested by means of the PWY and PSY supremum-ADF statistics. Following PSY, much of the literature maintains that, under the null hypothesis, the time series of interest behaves like a unit-root (i.e., integrated of order one) process with a local-to-zero drift that is dominated by the stochastic-trend component and vanishes as the length of the series grows beyond all bounds. We argue that the assumption of a weak, vanishing drift is far from innocuous and deviations from this assumption can result in severe over-rejection of the true unit-root hypothesis in favor of an explosive alternative. The effects of possible misspecification of the drift are quantified using Monte Carlo experiments under two different scenarios about the drift characteristics of the data, namely that of a stochastic drift that evolves as a covariance-stationary process and of a fixed drift the magnitude of which reflects estimates obtained from real-world time series (such as real stock prices). We also argue that using asymptotic critical values obtained under a DGP with dominant, non-vanishing drift is not without difficulties either since, unless the drift is substantial, right-tailed unit-root tests tend to under-reject for the sample sizes that are common in applications. Bootstrap-assisted versions of the PWY and PSY testing procedures, allowing for a drift comparable to that found in the observed data, are shown to offer an asymptotically valid – but only partially effective (in finitely-sized samples) – way of dealing with such difficulties. We consider a bootstrap-based technique to calibrate the significance level of tests, thus ensuring that the probability of erroneous rejections of the unit-root hypothesis is as close as possible to the desired target value.

The second issue relates to the dating of the origination and termination dates of multiple explosive episodes based on the PSY crossing principle. The latter amounts to comparing a sequence of recursively constructed ADF statistics to corresponding critical values (thresholds) obtained under a unit-root DGP. We argue that the standard practice of using conventional levels of significance for these critical values fails to guarantee control of the probability of detecting one or more explosive episodes when none are present in the data. Even though the origination and termination dates of (mildly) explosive episodes are consistently estimable, as long as the significance level associated with the relevant critical values approaches zero as the sample size becomes infinitely large, the undesirable tendency of the crossing rule to exaggerate the presence of explosiveness is an issue of concern when fixed significance levels are used. We demonstrate that the use of 5%-level critical values, for instance, in the PSY date-stamping procedure results in at least one explosive episode being detected in more than 70% of artificial samples from a unit-root DGP, the problem becoming more pronounced as the sample size increases. Such findings offer a partial explanation for the relatively large number of apparent explosive episodes identified in applied work that uses the PWY and PSY dating algorithms, and highlight the need for appropriate adjustments to be made in order to control the probability of false detections of explosiveness. We consider how this may be achieved by using a bootstrap-based technique to calibrate the significance level for the relevant critical values.

The importance of these practical considerations and the need for careful application of the PWY and PSY recursive techniques is further highlighted in our empirical study. Using monthly time series of U.S. real stock prices and real dividends for the period 1927:3–2020:6, we demonstrate that results relating to the detection of possible explosive episodes, and to the estimation of their origination and termination dates, can vary considerably depending on how the PSY testing and date-stamping procedures are implemented.

The remainder of the paper is organized as follows. Section 2 gives a brief description of the PWY and PSY testing and date-stamping recursive procedures. Sections 3 and 4 explore the properties of the PWY and PSY test procedures under data-generating mechanisms that involve a stochastic drift component and a non-vanishing drift of fixed

magnitude, respectively. Section 5 examines some properties of the PSY dating algorithm in the absence of explosive episodes in the data. Section 6 contains an analysis of the properties of U.S. stock prices and dividends. Section 7 summarizes and concludes.

2 The supremum-ADF approach

The procedures developed by PWY and PSY for detecting explosive behavior associated with rational bubbles rely on sequential implementation of right-tailed unit-root tests based on recursively constructed subsamples. In the absence of explosive episodes, the time series of interest $\{Y_t\}$ (e.g., asset prices) is assumed in PWY to have no deterministic drift, whereas PSY allow for a local-to-zero drift, in the sense that

$$Y_t = \delta T^{-\eta} + Y_{t-1} + u_t = \delta t T^{-\eta} + \sum_{j=1}^t u_j + Y_0, \quad (1)$$

for $t = 1, 2, \dots, T$, where δ is a non-zero constant and $\{u_t\}$ is a covariance-stationary process with mean zero (and spectral density that is bounded away from zero and infinity in a neighborhood of the origin). The localizing parameter $\eta \geq 0$ controls the behavior of the asymptotically negligible (as T approaches infinity) drift $\delta t T^{-\eta}$ of $\{Y_t\}$. This drift is dominated by the stochastic-trend component of $\{Y_t\}$ when $\eta > \frac{1}{2}$ (the latter being at most of order \sqrt{t} in probability) and is at least as strong as the stochastic trend when $\eta < \frac{1}{2}$. Phillips, Shi, and Yu (2014) argue that such a specification provides a reasonable description of the drift characteristics of many financial time series, and PSY focus on the case where $\eta > \frac{1}{2}$.

Unit-root tests are implemented in PWY and PSY using autoregressive models of the form

$$\Delta Y_t = \alpha + \theta Y_{t-1} + \sum_{i=1}^{\ell} \phi_i \Delta Y_{t-i} + \varepsilon_t, \quad (2)$$

for some integer $\ell \geq 0$, where Δ is the first-difference operator and $\{\varepsilon_t\}$ are white-noise errors, with the null and alternative hypotheses being $\theta = 0$ and $\theta > 0$, respectively.² Note that the inclusion of a deterministic linear trend as an additional covariate in (2) is not appealing in the context of right-tailed unit-root tests because it implies the presence

²As usual, the sum on the right-hand side of (2) is understood to be empty when $\ell = 0$.

of a deterministic component in $\{Y_t\}$ that has an empirically unrealistic form under the alternative hypothesis of a root greater than one (a point also made by Phillips, Shi, and Yu (2014)).

It is important to bear in mind that the PSY and PWY procedures have a dual goal: they provide a means of testing whether an explosive episode is present in a given sample; if the existence of such an episode is established, they also provide a way to estimate its origination and termination dates. Subsequent discussion will, therefore, focus on the properties of PWY and PSY tests, as well as on some aspects of the associated date-stamping algorithms.

2.1 The PWY procedure

The testing procedure of PWY is based on the supremum of the collection of ADF statistics computed over recursive subsamples the length of which increases by one observation at the time. More formally, for $0 \leq r_1 < r_2 \leq 1$, let $ADF(r_1, r_2)$ stand for the conventional least-squares t -statistic for testing the hypothesis $\theta = 0$ in (2) based on the subsample $\{Y_t; t = \lfloor Tr_1 \rfloor, \dots, \lfloor Tr_2 \rfloor\}$, where $\lfloor x \rfloor$ denotes the largest integer not exceeding the real number x . For some predetermined initial fraction $r_0 \in (0, 1)$ of the sample size T (PSY recommend setting $r_0 = 0.01 + 1.8T^{-1/2}$), the PWY test rejects for large values of the Supremum ADF (SADF) statistic

$$SADF(r_0) := \sup_{r_2 \in [r_0, 1]} ADF(0, r_2). \quad (3)$$

The asymptotic distribution of $SADF(r_0)$ under a no-drift ($\alpha = 0$) or localized-drift ($\alpha = \delta T^{-\eta}$) unit-root null hypothesis ($\theta = 0$) can be found in PWY and Phillips, Shi, and Yu (2014), respectively. In the presence of a local-to-zero drift, the asymptotic null distribution of $SADF(r_0)$ is not the same under $\eta > \frac{1}{2}$ and $\eta < \frac{1}{2}$.

When the unit-root null hypothesis is rejected in favor of the alternative that $\{Y_t\}$ exhibits explosive behavior in at least some parts of the sample, a simple crossing rule may be used to date-stamp the beginning and end of an explosive episode: the estimated origination (resp. termination) date of explosive behavior is the first observation for which $ADF(0, r_2)$ is sufficiently large (resp. small) relatively to the critical value associated with

a right-tailed unit-root test based on $ADF(0, r_2)$. More specifically, for some predetermined level of significance $\beta \in (0, 1)$, the estimated origination and termination dates of explosive behavior are $\lfloor T\hat{r}_e \rfloor$ and $\lfloor T\hat{r}_f \rfloor$, respectively, where

$$\begin{aligned}\hat{r}_e &:= \inf\{r_2 \in [r_0, 1] : ADF(0, r_2) > Q_\beta(r_2)\}, \\ \hat{r}_f &:= \inf\{r_2 \in [\hat{r}_e, 1] : ADF(0, r_2) < Q_\beta(r_2)\},\end{aligned}$$

and $Q_\beta(r_2)$ is the $(1 - \beta)$ -quantile of the distribution of $ADF(0, r_2)$ under the unit-root hypothesis. As discussed in PSY, it may be desirable to adjust the date-stamping algorithm so as to exclude short-lived explosive episodes (see next subsection).

2.2 The PSY procedure

The PWY procedure is generalized in PSY in an attempt to improve detection of multiple, periodically collapsing bubbles (e.g., such as those considered by [Evans \(1991\)](#)). The procedure is based on the supremum of a collection of ADF statistics computed over doubly recursive subsamples. More specifically, for some predetermined $r_0 \in (0, 1)$, the PSY test rejects for large values of the Generalized SADF statistic (GSADF) statistic

$$GSADF(r_0) := \sup_{\substack{r_1 \in [0, r_2 - r_0] \\ r_2 \in [r_0, 1]}} ADF(r_1, r_2) = \sup_{r_2 \in [r_0, 1]} BSADF(r_0, r_2), \quad (4)$$

where $BSADF(r_0, r_2)$ is the Backward SADF (BSADF) statistic defined as

$$BSADF(r_0, r_2) := \sup_{r_1 \in [0, r_2 - r_0]} ADF(r_1, r_2). \quad (5)$$

In other words, $BSADF(r_0, r_2)$ is obtained as the supremum of ADF statistics computed from recursive subsamples which are constructed by fixing the end observation $\lfloor Tr_2 \rfloor$ and then moving backwards in the sample, while $GSADF(r_0)$ is obtained as the double supremum of the ADF statistics computed over all possible recursive subsamples of minimum size $\lfloor Tr_0 \rfloor$. The asymptotic distributions of $GSADF(r_0)$ and $BSADF(r_0, r_2)$ under the localized-drift, unit-root null hypothesis ($\theta = 0$, $\alpha = \delta T^{-\eta}$, $\eta > \frac{1}{2}$) can be found in PSY.

In terms of date-stamping, the PSY rule is analogous to that of PWY. Specifically, for some predetermined $\beta \in (0, 1)$ and $\gamma > 0$, the estimated origination and termination

dates of explosive behavior are $\lfloor T\hat{r}_e \rfloor$ and $\lfloor T\hat{r}_f \rfloor$, respectively, where

$$\begin{aligned}\hat{r}_e &:= \inf\{r_2 \in [r_0, 1] : BSADF(r_0, r_2) > C_\beta(r_2)\}, \\ \hat{r}_f &:= \inf\{r_2 \in [\hat{r}_e + \gamma T^{-1} \ln T, 1] : BSADF(r_0, r_2) < C_\beta(r_2)\},\end{aligned}$$

and $C_\beta(r_2)$ is the $(1-\beta)$ -quantile of the distribution of $BSADF(r_0, r_2)$ under the unit-root hypothesis. The justification for the presence of the term $\gamma T^{-1} \ln T$ in the data-stamping rule is that, for an explosive episode to be meaningfully classified as a bubble, its duration should exceed a minimal period $\gamma \ln T$ (e.g., one year).

When multiple explosive episodes may be present in the data, the procedure can be used sequentially for date-stamping purposes. Specifically, after the termination date of the first explosive episode has been determined, criteria analogous to those above are used to estimate the location of a second such episode, conditionally on the first, and so on (see PSY and Phillips, Shi, and Yu (2015b) for more details). A similar sequential procedure may also be used in the context of the PWY approach, using the quantiles of the null sampling distribution of $ADF(0, r_2)$.

3 Stochastic drift and supremum-ADF tests

The PSY procedure relies heavily on the assumption that, under the null hypothesis of a unit root, the time series of interest has an asymptotically negligible drift that is dominated by its martingale component. To investigate how assumptions about the drift in the data affect the properties of supremum-ADF tests, we first consider the case where the DGP of the fundamental driving variable contains a drift component of a type that has been found to be empirically relevant.

3.1 Motivation

To motivate our analysis, we consider a simple present-value framework for the price of a stock and its dividend when the DGP for the latter has a time-varying drift. The intuition as to why such a DGP may pose additional difficulties when assessing possible explosive behavior in stock prices is simple. From the decomposition of asset returns due to

Campbell and Shiller (1989), the price–dividend ratio is known to depend on the expected growth rate of dividends and expected discount rates. Consequently, time-variation in the expected dividend growth rate will create an additional source of variation in stock prices. Moreover, it is well known that changes in the discount rate and/or the growth rate of dividends may generate explosive-looking behavior (see, inter alia, Driffill and Sola (1998) and Phillips and Yu (2011)).

Our framework is consistent with the empirical evidence presented in Pettenuzzo, Sabbatucci, and Timmermann (2020), who construct an econometric model that matches the features of their measure of dividend growth. Specifically, we focus on the component of their model which is found to be a strong predictor of dividend growth, namely a persistent component that captures a smoothly evolving time-varying mean.³ More formally, letting D_t denote the stock’s dividend payment between dates $t - 1$ and t , it is assumed that logarithmic dividends evolve according to the equations

$$\Delta \ln D_t = \mu_t + \sigma_d \varepsilon_t, \quad (6)$$

$$\mu_t - \mu = \phi(\mu_{t-1} - \mu) + \sigma_\mu \zeta_t, \quad (7)$$

where μ , ϕ , σ_d , and σ_μ are positive parameters (with $\phi < 1$). It is further assumed that $\{\varepsilon_t\}$ and $\{\zeta_t\}$ are sequences of independent, identically distributed (i.i.d.) $N(0, 1)$ random variables independent of each other. This implies that, conditional on information available at date t , $\ln D_t$ has drift μ_t , which evolves as a covariance-stationary autoregressive process with mean μ .

Under a standard no-arbitrage condition, the stock price at date t , denoted P_t , satisfies

$$P_t = e^{-\rho} \mathbf{E}_t(P_{t+1} + D_{t+1}), \quad (8)$$

where, for simplicity, the discount factor is specified as $e^{-\rho}$, $\rho > 0$ being the discount rate, and \mathbf{E}_t denotes conditional expectation given the information set available to market participants at date t . The solution to (8) is $P_t = P_t^F + P_t^B$, where $P_t^F = \sum_{j=1}^{\infty} e^{-\rho j} \mathbf{E}_t(D_{t+j})$

³Pettenuzzo, Sabbatucci, and Timmermann (2020) also include a jump component and a transitory component with time-varying volatility in their decomposition of dividend growth to capture specific features of the data, but we exclude such components since they are not essential in our context.

is the market-fundamentals component and $\{P_t^B\}$ is any sequence of random variables such that $P_t^B = e^{-\rho} \mathbf{E}_t(P_{t+1}^B)$. The component P_t^B is a rational bubble, reflecting the fact that an asset may have value because its price is expected to increase in future periods.

As shown in Appendix A.1, if the existence of bubbles can be ruled out in this framework (i.e., $P_t^B = 0$ for all t), stock prices are such that the price–dividend ratio $K_t := P_t/D_t$ satisfies

$$K_t = \sum_{j=1}^{\infty} \exp \left(j \left[\mu - \rho + \frac{\sigma_{\mu}^2}{2(1-\phi^2)} + \frac{\sigma_d^2}{2} \right] + \frac{\phi(1-\phi^j)(\mu_t - \mu)}{1-\phi} - \frac{\phi^2(1-\phi^{2j})\sigma_{\mu}^2}{2(1-\phi^2)^2} \right). \quad (9)$$

It is clear from (9) that the price–dividend ratio varies with the conditional mean μ_t of the dividend growth process. Therefore, a positive shock to μ_t that is observed by market participants will increase the price–dividend ratio. If ϕ is large, K_t will remain high for several periods, which may potentially be interpreted as a bubble by an external observer. Once μ_t reverts to its mean (or a negative shock to μ_t is observed), K_t will fall, which may be interpreted as a bubble burst.⁴

3.2 Simulation evidence

To assess how the presence of a time-varying drift in dividends affects results obtained by conventional implementation of the PWY and PSY testing procedures, we carry out a small Monte Carlo experiment based on artificial data generated according to (6), (7) and (9). The parameter values of the DGP, calibrated on the basis of the empirical results reported in [Pettenuzzo, Sabbatucci, and Timmermann \(2020\)](#), are: $\sigma_d = 0.0093$, $\mu = 0.0033$, $\phi = 0.443$, $\sigma_{\mu} = 0.0155$, $\rho = 0.004732$. Details on how the parameters are calibrated are provided in Appendix A.2.

To reflect what is common practice in applied work, we focus on logarithmic prices, generated as $\ln P_t = \ln K_t + \ln D_t$. In each of 1000 Monte Carlo replications, we compute the values of the SADF and GSADF statistics defined in (3) and (4), and compare them

⁴Note that the setting considered here is different from that of [Phillips and Shi \(2019\)](#), in whose model of stock-market crashes the drift in the logarithmic price–dividend ratio is inherited from a two-regime DGP for logarithmic prices allowing for a local-to-zero drift in one regime and a stochastic drift in the form of i.i.d. uniform random variables in the other (with logarithmic dividends being a random walk with time-invariant drift).

T	SADF			GSADF		
	0.10	0.05	0.01	0.10	0.05	0.01
Asymptotic Critical Values						
100	0.127	0.082	0.0277	0.149	0.104	0.034
200	0.212	0.123	0.028	0.211	0.130	0.031
400	0.282	0.161	0.035	0.240	0.148	0.049
800	0.403	0.266	0.077	0.403	0.260	0.096
Finite-Sample Critical Values						
100	0.160	0.091	0.0230	0.159	0.085	0.019
200	0.212	0.123	0.044	0.188	0.112	0.013
400	0.283	0.161	0.036	0.265	0.145	0.030
800	0.385	0.256	0.082	0.335	0.191	0.057

Table 1: Rejection frequencies of recursive unit-root tests for logarithmic prices

with the asymptotic and finite-sample critical values given in Table 1 of PSY. Recall that these critical values are obtained under the assumption that the data satisfy a random-walk model with asymptotically vanishing drift $1/T$. The proportion of Monte Carlo replications in which the null hypothesis of a unit root in $\ln P_t$ is rejected in favor of an explosive alternative (at standard levels of significance) is reported in Table 1. (Unless the value of ℓ in the autoregressive model (2) that is used to construct ADF-type statistics is explicitly stated, it should henceforth be understood to be zero. Similarly, the initialization parameter r_0 used in the implementation of recursive procedures is always to be understood to be $r_0 = 0.01 + 1.8T^{-1/2}$.)

It is immediately obvious that the test procedures reject the unit-root hypothesis too frequently. Although logarithmic dividends are not explosive and the price–dividend ratio is covariance-stationary, the two tests detect explosiveness in logarithmic prices in a large proportion of the artificial samples. This over-rejection of the null hypothesis can be attributed to two main reasons. First, as noted in Section 3.1, changes in the expected dividend growth result in changes in the price–dividend ratio, which could be mistaken for explosive behavior. This observation is obviously not new – as pointed out by [Driffill and Sola \(1998\)](#), among others, changes in fundamentals can generate bubble-like behavior in the data. It is worth noting, however, that the drift process is not very persistent in our

experiments; the value of ϕ is less than 0.45, so the effects of shocks to μ_t tend to dissipate rather quickly. Nevertheless, the SADF and GSADF tests reject in favor of explosive behavior in prices far too often. The second reason for our findings could be the underlying specification of the null model. Under our DGP, $\ln D_t$ is a random walk with a stochastic drift whose mean is non-zero, and such characteristics are inherited by $\ln P_t$. However, the critical values used to compute the rejection frequencies of the SADF and GSADF tests in Table 1 are obtained under a unit-root DGP having an asymptotically vanishing drift that is dominated by the stochastic trend. The latter assumption is extremely common in applications involving the PWY and PSY procedures, but is far from innocuous, especially in view of the fact that many financial series (including time series such as dividends and stock prices) tend to exhibit a positive drift the magnitude of which does not decrease as the length of the series increases.

The primary justification for allowing for a local-to-zero drift under the null hypothesis is the claim that, for the sample sizes typically encountered in applications, the drift effect is dominated by the stochastic trend associated with the martingale component of the data. Thus, the use of critical values for unit-root tests obtained under a DGP with a non-negligible drift would be misleading in finite samples. We will argue in the sequel that, for sample sizes and parameters values that closely resemble those of financial series such as stock prices, the rejection probabilities of unit-root tests are quite sensitive with respect to the magnitude of the drift and the volatility of the shocks. Using critical values for the tests that are based on the assumption of a weak, local-to-zero drift ($\eta = 1$) can result in tests that are substantially liberal (in that the actual rejection probability under the null hypothesis is much larger than the nominal level). Equally, in samples of sizes that are typical in applications, the use of asymptotic critical values based on the assumption of a dominating drift of fixed magnitude ($\eta = 0$) can yield tests that are severely conservative (in that the actual rejection probability under the null hypothesis is much smaller than the nominal level). We turn our attention to these issues next.

4 Non-vanishing drift and supremum-ADF tests

In this section, we explore further the properties of the PWY and PSY test procedures in the presence of a realistically-sized, time-invariant drift in the DGP.

4.1 Misspecification of drift characteristics

Many financial time series, including stock prices and dividends, appear to be described well as unit-root processes (either in levels or in logarithms) with drift of a substantial magnitude. However, as is well known from Phillips, Shi, and Yu (2014), under a DGP such as (1), or (2) with $\theta = 0$, the asymptotic behavior of ADF statistics varies depending on whether $0 \leq \eta < \frac{1}{2}$ or $\eta > \frac{1}{2}$. Specifically, critical values for a right-tailed unit-root test under $\eta = 0$ are larger than those obtained under $\eta > \frac{1}{2}$. Phillips, Shi, and Yu (2014) regard an autoregressive process such as (2) with $\theta > 0$, $\alpha = \delta T^{-\eta}$ and $\eta = 0$ as empirically unrealistic because the coexistence of a non-zero intercept and a root greater than one generates a dominating deterministic component with an explosive form. They recommend that in applied work results should be presented for a range of values of η . However, most empirical applications of right-tailed unit-root tests rely on asymptotic critical values obtained under $\eta > \frac{1}{2}$ (that is, under a unit-root null where the drift is not the dominant characteristic and vanishes asymptotically), or use simulated finite-sample critical values obtained under the same assumption. Therefore, if the DGP is a unit-root process with non-vanishing drift (which is plausible for many financial series), it is likely that the unit-root hypothesis will be wrongly rejected in favor of an explosive alternative more frequently than the nominal level of tests implies. This was indeed found to be the case under the time-varying dividend-drift scenario considered in Section 3.

Focusing on the statistics defined in (3) and (4), it is known from Phillips, Shi, and Yu (2014) that, under $\theta = 0$, $\alpha = \delta T^{-\eta}$ and $\eta = 0$ in (2),

$$SADF(r_0) \rightsquigarrow \sup_{r \in [r_0, 1]} \left\{ \frac{\sqrt{3}}{r^{3/2}} \left[\int_0^r s dW(s) - \int_0^r W(s) ds \right] \right\}, \quad (10)$$

as $T \rightarrow \infty$, where $\{W(s)\}$ is a standard Brownian motion on $[0, 1]$ and “ \rightsquigarrow ” denotes

	SADF			GSADF		
	0.90	0.95	0.99	0.90	0.95	0.99
$r_0 = 0.190$	2.4908	2.7917	3.3097	3.6329	3.8710	4.3291
$r_0 = 0.137$	2.5428	2.8342	3.3648	3.7591	4.0234	4.4445
$r_0 = 0.100$	2.5486	2.8184	3.3995	3.8884	4.0976	4.4533
$r_0 = 0.074$	2.6319	2.9208	3.5127	3.9862	4.1836	4.5856
$r_0 = 0.055$	2.6837	2.9788	3.5626	4.0862	4.3308	4.7760

Table 2: Quantiles of the asymptotic distributions of the SADF and GSADF statistics when $\eta = 0$

convergence in distribution. Furthermore, as shown in Appendix A.3 (see Proposition A.1),

$$GSADF(r_0) \rightsquigarrow \sup_{\substack{r_1 \in [0, r_2 - r_0] \\ r_2 \in [r_0, 1]}} \left\{ \frac{\sqrt{3}}{(r_2 - r_1)^{3/2}} \left[(r_2 - r_1) \{W(r_2) + W(r_1)\} - 2 \int_{r_1}^{r_2} W(s) ds \right] \right\}, \quad (11)$$

as $T \rightarrow \infty$. Selected quantiles of the distributions of the limit random variables in (10) and (11), for the values of r_0 recommended in PSY, are given in Table 2.⁵

To quantify the implications of misspecification of the drift, we carry out Monte Carlo experiments. In order to ensure comparability with PSY, artificial stock prices are simulated according to a simple present-value model in which logarithmic dividends follow a Gaussian random walk with drift, generated as

$$\Delta \ln D_t = \mu + \sigma_d \varepsilon_t, \quad t \geq 1, \quad (12)$$

where $\{\varepsilon_t\}$ are i.i.d. $N(0, 1)$ random variables. Prices are determined via the condition

$$P_t = \rho^{-1} \mathbf{E}_t(P_{t+1} + D_{t+1}), \quad (13)$$

for some fixed discount rate $\rho > 0$, so that the market-fundamentals component of P_t is

$$P_t^F = \frac{\rho \exp(\mu + \frac{1}{2}\sigma_d^2)}{1 - \rho \exp(\mu + \frac{1}{2}\sigma_d^2)} D_t. \quad (14)$$

Since we are interested in the performance of tests in the absence of bubbles, we set $P_t = P_t^F$ for all t . The values of the DGP parameters are taken to be the same as in

⁵These are obtained via numerical simulation of the functionals in (10) and (11), using 2000 Monte Carlo replications, approximating the Brownian motions using partial sums of independent $N(0, 1)$ random variates with 1000 steps.

T	SADF			GSADF		
	0.10	0.05	0.01	0.10	0.05	0.01
	Asymptotic Critical Values					
100	0.126	0.073	0.024	0.144	0.091	0.036
200	0.155	0.087	0.019	0.185	0.103	0.030
400	0.265	0.166	0.046	0.241	0.157	0.038
800	0.355	0.210	0.065	0.352	0.245	0.074
1600	0.475	0.317	0.108	0.518	0.345	0.129
	Finite-Sample Critical Values					
100	0.162	0.083	0.019	0.151	0.072	0.027
200	0.155	0.089	0.026	0.169	0.084	0.014
400	0.269	0.166	0.047	0.262	0.153	0.025
800	0.360	0.211	0.064	0.287	0.160	0.044
1600	0.443	0.291	0.083	0.421	0.260	0.082

Table 3: Rejection frequencies of recursive unit-root tests for logarithmic prices (assuming $\eta = 1$)

one of the Monte Carlo designs in PSY: $\rho = 0.985$, $\mu = 0.001$, $\sigma_d = 0.01$, $D_0 = e$; we will henceforth refer to this set of parameter values as the ‘baseline configuration’.⁶ As in the simulations in Section 3.2, we focus on logarithmic prices to reflect what is common practice in applied work (PWY and Phillips, Shi, and Yu (2014) also consider logarithmic prices in their empirical illustrations). For each of 1000 artificial samples, we compute the values of the SADF and GSADF statistics for $\ln P_t$ and compare them with the asymptotic and finite-sample critical values given in Table 1 of PSY, which correspond to the case of a misspecified drift ($\eta = 1$) under our DGP. The proportion of samples in which the unit-root hypothesis is rejected in favor of an explosive alternative (at standard levels of significance) is reported in Table 3.

The SADF and GSADF tests over-reject in all cases, the deviation of the estimated rejection probabilities of the tests from the corresponding nominal level increasing as the sample size grows. This is the result of using critical values for the tests based on the assumption that the DGP is a random walk with a mild and asymptotically negligible

⁶The baseline configuration is based on Robert Shiller’s monthly time series of logarithmic real dividends for the S&P 500 stocks.

T	SADF			GSADF		
	0.10	0.05	0.01	0.10	0.05	0.01
100	0.005	0.003	0.001	0.001	0.001	0.001
200	0.005	0.003	0.000	0.003	0.002	0.000
400	0.004	0.002	0.001	0.001	0.000	0.000
800	0.007	0.003	0.000	0.000	0.000	0.000
1600	0.017	0.005	0.001	0.000	0.000	0.000

Table 4: Rejection frequencies of recursive unit-root tests for logarithmic prices (assuming $\eta = 0$)

drift when the true drift is dominant and does not vanish as the length of the simulated time series increases. It would seem, therefore, that for the sample sizes and parameter values that are commonly encountered in applications, the assumption of a mild, vanishing drift is not innocuous and deviations from this assumption can have a deleterious effect on the level accuracy of SADF and GSADF tests.

The alternative approach of using critical values associated with the asymptotic results (10) and (11) is not without difficulties either. Table 4 contains the Monte Carlo rejection frequencies of the SADF and GSADF tests, under the same design as before, when critical values from Table 2, obtained under the assumption that $\eta = 0$, are used. Even though the drift is now correctly specified, the unit-root null hypothesis is very rarely rejected. This is the opposite of what happens when critical values obtained under $\eta = 1$ are used. Additional experiments (not reported here) reveal that the severe under-rejection of the unit-root hypothesis is due to the poor quality of asymptotic approximations to the null distributions of the SADF and GSADF statistics for relatively small values of the drift. When the drift is more pronounced (e.g., $\mu = 0.5$), although admittedly less empirically plausible, the tests tend to have rejection probabilities which are close to the nominal level for the sample sizes considered here.

4.2 Robustness checks

To assess the robustness, or lack thereof, of our findings with respect to different values of the parameters, we carry out additional simulation experiments, using (12) and (14) as

the DGP and setting $P_t = P_t^F$ for all t . Specifically, we repeat the earlier Monte Carlo experiments, varying the value of one of the DGP parameters at the time while keeping all others fixed at the values associated with the baseline configuration. The values of μ and σ_d considered are: $\mu \in \{0.0005, 0.001, 0.002, 0.003, 0.004\}$, $\sigma_d \in \{0.005, 0.01, 0.015, 0.02\}$. The results of the simulations are collected in Appendix A.4.

When critical regions for the SADF and GSADF tests are constructed under the assumption of a local-to-zero drift with localizing parameter $\eta = 1$, differences between the simulation-estimated and nominal levels of the tests become more pronounced as the drift parameter μ increases. This is not surprising since the larger μ is, the more dominant the deterministic time trend is over the stochastic trend in the data, and the quality of asymptotic approximations to the distributions of the test statistics obtained under the assumption of a mild and asymptotically vanishing drift suffers as a result. The reverse is true for the effect of the variance parameter σ_d on the rejection probabilities of the tests, the latter being closer to the nominal level the larger σ_d is. This is not an unexpected finding since the relative importance of the stochastic trend relative to a linear trend increases the larger the variance of the noise is. As a result, approximations to the null distributions of the SADF and GSADF statistics obtained under the assumption that the order of magnitude of the data is that of a random walk (i.e., $\eta > \frac{1}{2}$) are likely to be more accurate the larger σ_d is.

In summary, the overall message from the simulation exercises in Sections 4.1 and 4.2 (as well as those in Section 3.2) is that, for empirically relevant parameter values and sample sizes, the possibility of a non-vanishing drift cannot be safely ignored. Furthermore, the difference between the actual probability that the tests incorrectly reject the null hypothesis and the desired (nominal) probability is heavily dependent on the values of the parameters of the DGP and on the size of the sample. Large samples, large drift values, and small volatility all exacerbate the problem of over-rejection of the unit-root hypothesis observed under an erroneous assumption of a drift lying in a neighborhood of zero.⁷

⁷Wang and Yu (2022) provide evidence that a GSADF test relying on a first-order autoregressive model and the PSY critical values also tends to over-reject in favor of an explosive alternative under a DGP different from those considered here, namely one which involves an one-off shift between a driftless unit-root process and a process that is covariance-stationary about a deterministic quadratic time trend.

4.3 Bootstrap-assisted tests

Instead of using the PWY/PSY vanishing-drift critical values or those associated with the fixed-drift results (10) and (11), one may rely on critical values for the SADF and GSADF tests obtained by means of a suitable bootstrap procedure. [Etienne, Irwin, and Garcia \(2014\)](#), [Harvey, Leybourne, Sollis, and Taylor \(2016\)](#), [Phillips and Shi \(2019\)](#), [Phillips and Shi \(2020\)](#), and [Monschang and Wilfing \(2021\)](#), for example, use the wild bootstrap in the context of the PSY and/or PWY testing procedures, restricting the bootstrap data to have no drift, while [Hafner \(2020\)](#) allows for a fixed drift in the bootstrap DGP used to implement the SADF test.

We consider here a somewhat similar bootstrap procedure for tests based on the autoregressive model (2), ensuring that bootstrap data have a drift which reflects the characteristics of the observed data. To describe the procedure, let $(\tilde{\alpha}, \tilde{\phi}_1, \dots, \tilde{\phi}_\ell)$ be least-squares estimates of the coefficients of (2) under the constraint $\theta = 0$ and $\{\tilde{\varepsilon}_t\}$ be the corresponding residuals. Our wild-bootstrap scheme amounts to generating a bootstrap replicate $\{Y_t^*\}$ of the observable time series $\{Y_t\}$ according to

$$Y_t^* = \tilde{\alpha} + Y_{t-1}^* + \sum_{i=1}^{\ell} \tilde{\phi}_i \Delta Y_{t-i}^* + \nu_t^* \tilde{\varepsilon}_t, \quad t = \ell + 1, \ell + 2, \dots, T, \quad (15)$$

with $Y_t^* = Y_t$ for $t \leq \ell$ and $\{\nu_t^*\}$ being i.i.d. random variables, independent of $\{Y_t\}$, having mean zero and variance one.⁸ For a sufficiently large integer $B > 0$ and any $\beta \in (0, 1)$, β -level bootstrap critical values for the SADF and GSADF tests may then be obtained as the $\lfloor (B+1)(1-\beta) \rfloor$ -th largest of the values of bootstrap replicates of the $SADF(r_0)$ and $GSADF(r_0)$ statistics, respectively, computed from B independent (conditionally on the original data) realizations of $\{Y_t^*\}$ from the bootstrap DGP (15). In the implementation of the wild bootstrap in the remainder of the paper we take $\{\nu_t^*\}$ to be $N(0, 1)$ random variables, but other choices are also possible.⁹

⁸The wild bootstrap allows for the possibility of conditional heteroskedasticity in the errors in (2). If this is not a concern, the ordinary bootstrap may be used instead, which amounts to replacing $\nu_t^* \tilde{\varepsilon}_t$ in (15) with bootstrap errors that are chosen by sampling randomly, with replacement, from the residuals $\{\tilde{\varepsilon}_t\}$. [Gutierrez \(2013\)](#) uses the latter resampling scheme, restricting the bootstrap DGP to have no drift.

⁹Popular alternatives include discretely distributed random variables taking the values ± 1 with equal probability, or the values $(1 \pm \sqrt{5})/2$ with probability $(\sqrt{5} \mp 1)/(2\sqrt{5})$, respectively.

T	SADF			GSADF		
	0.10	0.05	0.01	0.10	0.05	0.01
100	0.033	0.016	0.001	0.030	0.009	0.000
200	0.067	0.020	0.001	0.027	0.009	0.000
400	0.081	0.036	0.003	0.032	0.013	0.001
800	0.093	0.042	0.005	0.038	0.015	0.001
1600	0.106	0.060	0.008	0.076	0.028	0.002

Table 5: Rejection frequencies of bootstrap-assisted recursive unit-root tests for logarithmic prices

As shown in Appendix A.3 (see Proposition A.2 and the ensuing discussion), the wild bootstrap provides consistent estimators of the null distributions of the SADF and GSADF statistics in the presence of a non-vanishing drift.¹⁰ This implies that SADF and GSADF tests constructed using wild-bootstrap critical values are asymptotically correct, in the sense that, under the unit-root hypothesis, their rejection probabilities converge to the nominal level β as T and B tend to infinity.

To get some insight into the finite-sample properties of bootstrap-assisted SADF and GSADF tests, we carry out simulation experiments, using (12) and (14) as the DGP and setting $P_t = P_t^F$ for all t . The parameter values used are those associated with our baseline configuration, so results are comparable with those reported in Section 4.1. The proportion of 1000 Monte Carlo replications in which the SADF and GSADF tests reject the unit-root hypothesis for $\ln P_t$ (at standard levels of significance), when wild-bootstrap critical values (with $B = 1000$) are used, is reported in Table 5.

The unit-root hypothesis tends to be rejected in favor of an explosive alternative less frequently than the nominal level of the tests implies, under-rejection being more severe in the case of the GSADF test. In fact, the bootstrap-assisted version of the SADF test has estimated rejection probabilities which are generally close to the nominal level when the sample size is relatively large, and is a clear improvement over SADF tests that rely on vanishing-drift or fixed-drift asymptotic critical values, as comparison with the results in Tables 3 and 4 reveals.

¹⁰Bootstrap consistency also holds when there is no drift under the null hypothesis.

Additional simulation experiments (not reported here) show that using the true – albeit unavailable in real-world applications – drift $\mu = 0.001$ in place of the estimated drift $\tilde{\alpha} = T^{-1} \sum_{t=1}^T \Delta \ln P_t$ to generate bootstrap replicates of the data does not improve the level accuracy of the GSADF test significantly, the test continuing to reject too infrequently even for the largest of the sample sizes considered. There is improvement, however, in the level accuracy of the SADF test, suggesting that the use of an estimated drift is not without cost.

In summary, although the use of the wild bootstrap in the context of the PSY and PWY testing procedures is to a certain extent beneficial when there is a dominant drift in the observed data, it may not guarantee effective control of the finite-sample errors in the level of recursive right-tailed unit-root tests. The bootstrap-assisted GSADF test seems to be particularly problematic in this respect, having a tendency to reject the unit-root hypothesis too rarely.^{11,12}

4.4 Calibrated bootstrap-assisted tests

The severity of the difficulties identified in the preceding section relating to the level accuracy of wild-bootstrap tests in the presence of a non-vanishing drift can potentially be mitigated by using a suitable bootstrap procedure to calibrate the nominal significance level of the tests so that their (estimated) finite-sample rejection probabilities under the unit-root hypothesis are approximately equal to some desired nominal value. Bootstrap-based calibration, as a method for improving the accuracy of approximate inferential procedures via level adjustments, dates back to [Loh \(1987\)](#); when applied to bootstrap-

¹¹In simulation experiments in which artificial data have a vanishing local-to-zero drift and their bootstrap replicates are driftless, [Monschang and Wilfing \(2021\)](#) also find SADF and GSAF tests that rely on wild-bootstrap critical values to be too conservative.

¹²It is worth noting that replacing $\nu_t^* \tilde{\varepsilon}_t$ in (15) with bootstrap errors that are drawn randomly, with replacement, from $\{\tilde{\varepsilon}_t\}$ results in bootstrap-assisted tests which exhibit small level distortions for time series which are not too short. This can be seen in [Table A8](#) (top panel) in [Appendix A.5](#), and suggests that the localized selection of bootstrap errors that the wild-bootstrap scheme involves has a deleterious effect on level accuracy under the DGP used in our experiments, which exhibits no conditional heteroskedasticity. The ordinary-bootstrap scheme is valid under such a DGP, but it would not necessarily be appropriate in heteroskedastic settings.

assisted tests, as is the case here, it amounts to the prepivoting method of [Beran \(1988\)](#).¹³

For a given nominal level $\beta \in (0, 1)$ and sufficiently large positive integers B and N , the calibration procedure for the bootstrap-assisted GSADF test consists of the following steps (the procedures for SADF is entirely analogous):

- (i) construct B bootstrap time series $\{Y_{b,t}^*\}$, $b = 1, 2, \dots, B$, of the same length as $\{Y_t\}$, using the wild-bootstrap scheme described in [Section 4.3](#);
- (ii) compute a collection $\{GSADF_b^*(r_0); b = 1, 2, \dots, B\}$ of (first-round) bootstrap GSADF values by applying the definition of $GSADF(r_0)$ to each $\{Y_{b,t}^*\}$ in place of $\{Y_t\}$;
- (iii) for each $\{Y_{b,t}^*\}$, construct N further bootstrap time series $\{Y_{b,i,t}^{**}\}$, $i = 1, 2, \dots, N$, of the same length as $\{Y_{b,t}^*\}$, by applying the wild-bootstrap scheme to $\{Y_{b,t}^*\}$ in place of $\{Y_t\}$;
- (iv) for each $b = 1, 2, \dots, B$, compute a collection $\{GSADF_{b,i}^{**}(r_0); i = 1, 2, \dots, N\}$ of (second-round) bootstrap GSADF values by applying the definition of $GSADF(r_0)$ to the (b, i) -th bootstrap time series $\{Y_{b,i,t}^{**}\}$ in place of $\{Y_t\}$;
- (v) for a grid of values of $\lambda \in (0, 1)$ (in a neighborhood of β) compute the proportion $\pi^*(\lambda)$ of $\{GSADF_b^*(r_0); b = 1, 2, \dots, B\}$ which exceed the $\lfloor (N+1)(1-\lambda) \rfloor$ -th largest of $\{GSADF_{b,i}^{**}(r_0); i = 1, 2, \dots, N\}$;
- (vi) find the value $\tilde{\lambda}$ for which $\pi^*(\tilde{\lambda}) = \beta$;¹⁴
- (vii) use the $\lfloor (B+1)(1-\tilde{\lambda}) \rfloor$ -th largest of $\{GSADF_b^*(r_0); b = 1, 2, \dots, B\}$ as the calibrated β -level bootstrap critical value for $GSADF(r_0)$.

To assess the efficacy of the bootstrap-based calibration procedure, we carry out simulation experiments, using [\(12\)](#) and [\(14\)](#) as the DGP and setting $P_t = P_t^F$ for all t . The parameter values used are once again those associated with our baseline configuration, so results are comparable with those reported in [Sections 4.1](#) and [4.3](#). The proportion of 1000

¹³Calibrating an asymptotically correct bootstrap procedure results in an asymptotically correct procedure again, but typically one with approximation errors of a smaller order of magnitude.

¹⁴The value $\tilde{\lambda}$ may be found by linearly interpolating $\pi^*(\lambda)$ between the grid values of λ .

T	SADF			GSADF		
	0.10	0.05	0.01	0.10	0.05	0.01
100	0.089	0.039	0.010	0.062	0.033	0.006
200	0.106	0.054	0.009	0.076	0.028	0.005
400	0.109	0.059	0.010	0.098	0.037	0.007
800	0.105	0.055	0.009	0.089	0.043	0.007
1600	0.108	0.069	0.012	0.129	0.068	0.012

Table 6: Rejection frequencies of calibrated bootstrap-assisted recursive unit-root tests for logarithmic prices

Monte Carlo replications in which SADF and GSADF tests reject the unit-root hypothesis for $\ln P_t$ (at standard levels of significance), when calibrated wild-bootstrap critical values (with $B = 1000$ and $N = 500$) are used, is reported in Table 6.

The results show that bootstrap calibration offers an effective way of reducing errors in the rejection probabilities of tests based on wild-bootstrap critical values when a non-vanishing drift is present under the unit-root hypothesis. Tests relying on calibrated bootstrap critical values have simulated rejection probabilities which are much closer to the nominal level compared with their non-calibrated counterparts or with tests that rely on asymptotic critical values. As with non-calibrated bootstrap-assisted tests, the calibrated SADF test fares somewhat better than the GSADF test in terms of finite-sample level accuracy, the former being on target in the majority of cases.¹⁵ Based on this evidence, the calibration method, though computationally intensive, has much to recommend it, and its use in the context of the PWY and PSY testing procedures is therefore advisable.

5 Non-explosiveness and date-stamping

In this section, we highlight and explore a potential difficulty associated with the practical implementation of the PSY date-stamping algorithm.

¹⁵Bootstrap calibration also offers level-accuracy improvements in the case of tests based on ordinary-bootstrap critical values, as the simulation results in Table A8 (bottom panel) in Appendix A.5 show.

5.1 Backward supremum-ADF tests

As noted in Section 2.2, the PSY procedure for estimating the origination and termination dates of explosive episodes is based on a simple crossing rule involving the sequence of backward SADF statistics defined in (5): if the value of the BSADF statistic exceeds a certain threshold (critical value), a period of explosiveness is estimated to start; when the value of BSADF falls below the threshold, the explosive period is estimated to end. Requiring the significance level β associated with these thresholds to tend to zero as the sample size T goes to infinity ensures that the critical values $C_\beta(r_2)$ used in the crossing rule grow beyond all bounds, and thus the probability of incorrectly detecting an explosive episode is asymptotically zero when data are generated by a DGP such as (1). Furthermore, Phillips, Shi, and Yu (2015b) show that the PSY procedure provides consistent estimates of the origination and termination dates of multiple mildly explosive episodes,¹⁶ as long as $C_\beta(r_2) \rightarrow \infty$ as $T \rightarrow \infty$ at a suitably slow rate. The same is true for the PWY procedure, in the presence of a single explosive episode, provided β is chosen as a function of T so that $Q_\beta(r_2) \rightarrow \infty$ sufficiently slowly as $T \rightarrow \infty$. A key point, however, is how the sequence of critical values relevant to this rule is constructed and used in practice.

In a typical implementation of the PSY procedure, a sufficiently large number M of independent artificial realizations of $\{Y_t\}$ are generated according to

$$\Delta Y_t = [Tr_2]^{-1} + u_t, \quad t = 1, 2, \dots, T,$$

for each $r_2 \in \{r_0, T^{-1}(\lfloor Tr_0 \rfloor + 1), T^{-1}(\lfloor Tr_0 \rfloor + 2), \dots, 1\} =: \mathcal{R}_2$, with $r_0 = 0.01 + 1.8T^{-1/2}$, $Y_0 = 0$, and $\{u_t\}$ being i.i.d. $N(0, 1)$ random variates. For each $r_2 \in \mathcal{R}_2$, these realizations are used to compute collections $\{BSADF_i(r_0, r_2); i = 1, 2, \dots, M\}$ of BSADF values and, for any $\beta \in (0, 1)$, an estimate $\hat{C}_\beta(r_2)$ of the $(1 - \beta)$ -quantile of the sampling distribution of $BSADF(r_0, r_2)$ is then obtained as the $\lfloor (M + 1)(1 - \beta) \rfloor$ -th largest of the values $BSADF_1(r_0, r_2), \dots, BSADF_M(r_0, r_2)$. The collection of simulation-estimated quantiles $\{\hat{C}_\beta(r_2); r_2 \in \mathcal{R}_2\}$ is used for the date-stamping of explosive episodes in the manner discussed in Section 2.2.

¹⁶These are characterized by the presence of an autoregressive root that is explosive in a neighborhood of unity, that is, of the form $1 + cT^{-a}$ for some $c > 0$ and $a \in (0, 1)$.

As already noted, the consistency results of PSY and Phillips, Shi, and Yu (2015b) require that the nominal significance level β for individual BSADF tests decreases to zero at an appropriate rate as the sample size increases, thus ensuring that, under the unit-root null hypothesis, the probability of detecting an explosive episode approaches zero asymptotically. However, as PSY (p. 1052) remark: “In empirical applications, it is also often convenient to fix β at some predetermined level such as 0.05 instead of using a drifting significance level”; PWY (p. 207) also point out that: “In practice, it is conventional to set the significance level in the 1–5% range.” This practice has, in fact, become the standard way of implementing the PSY procedure in applied work. Not unexpectedly perhaps, such an approach is not without difficulties, particularly when the whole sequence of BSADF statistics involved in the procedure is considered, as is the case when dating multiple explosive episodes.

To explain, recall that, in order to estimate the origination and termination dates of multiple explosive episodes in an observed time series $\{Y_t\}$ of length T , the entire collection of statistics $\{BSADF(r_0, r_2); r_2 \in \mathcal{R}_2\}$ needs to be considered, with an explosive episode deemed to begin (resp. end) when $BSADF(r_0, r_2)$ is greater (resp. smaller) than the corresponding critical value $\hat{C}_\beta(r_2)$. If $\{Y_t\}$ satisfies (1) with $\eta > \frac{1}{2}$, the method described above for obtaining $\{\hat{C}_\beta(r_2); r_2 \in \mathcal{R}_2\}$ ensures that, for large enough T and M , the probability of $BSADF(r_0, r_2)$ exceeding $\hat{C}_\beta(r_2)$ for any given $r_2 \in \mathcal{R}_2$ is approximately equal to β . However, if each of the $BSADF(r_0, r_2)$ statistics involved in the dating procedure is used in conjunction with a critical value of fixed nominal level $\beta \in (0, 1)$, the probability of $BSADF(r_0, r_2)$ exceeding $\hat{C}_\beta(r_2)$ for some $r_2 \in \mathcal{R}_2$ when no explosive episodes are present can evidently be much greater than β .¹⁷ In other words, in the absence of explosiveness, the probability of detecting at least one explosive episode using the crossing

¹⁷Defining $\mathcal{Q}_\beta := \{r_2 \in \mathcal{R}_2 : BSADF(r_0, r_2) > \hat{C}_\beta(r_2)\}$ and letting $|\mathcal{A}|$ denote the cardinality of the set \mathcal{A} , it follows by the inequality due to Kuai, Alajaji, and Takahara (2000) that, in the absence of explosiveness,

$$\mathbb{P}(|\mathcal{Q}_\beta| \geq 1) = \mathbb{P}\left(\bigcup_{r_2 \in \mathcal{R}_2} \{BSADF(r_0, r_2) > \hat{C}_\beta(r_2)\}\right) \geq \beta \sum_{r_2 \in \mathcal{R}_2} \left(\frac{1}{\lfloor S_{r_2} \rfloor} - \frac{S_{r_2} - \lfloor S_{r_2} \rfloor}{(1 + \lfloor S_{r_2} \rfloor)\lfloor S_{r_2} \rfloor}\right),$$

where $S_{r_2} := \sum_{j \in \mathcal{R}_2} \mathbb{P}[BSADF(r_0, j) > \hat{C}_\beta(j) | BSADF(r_0, r_2) > \hat{C}_\beta(r_2)]$. If $\{BSADF(r_0, r_2); r_2 \in \mathcal{R}_2\}$ were (asymptotically) independent under (1), then $\mathbb{P}(|\mathcal{Q}_\beta| \geq 1) \approx 1 - (1 - \beta)^{|\mathcal{R}_2|}$.

principle (or, equivalently, the expected proportion of originating explosive episodes that are erroneously identified as such by the crossing principle) can be substantially larger than what is typically thought of as the nominal level of individual BSADF tests. Unless $\hat{C}_\beta(r_2)$ increases to infinity with T , the problem of explosive episodes being erroneously detected with high probability remains when such episodes are considered to be genuine only when $BSADF(r_0, r_2)$ exceeds $\hat{C}_\beta(r_2)$ for several consecutive values of $r_2 \in \mathcal{R}_2$, as recommended by PSY.¹⁸

5.2 Numerical results

To illustrate numerically the difficulties discussed in the previous subsection, we generate 1000 artificial time series satisfying (1), with $\delta = \eta = 1$ and $\{u_t\}$ being i.i.d. $N(0, 1)$ variates, and compute the values of the statistics $\{BSADF(r_0, r_2); r_2 \in \mathcal{R}_2\}$ in each case. The proportion of artificial time series for which the BSADF statistics exceed, at least once, the relevant 0.05-level critical values for a minimum of $k \in \{1, 2, 3, 4, 5, 6, 9, 12\}$ consecutive values of r_2 are shown in Table 7 (top panel).¹⁹ In addition, we compute the average number of explosive episodes per time series detected by the PSY crossing rule, and record the results in Table 7 (bottom panel). An explosive episode is defined as the event that the BSADF sequence exceeds the relevant 0.05-level critical values for at least k consecutive periods; the episode is considered to have terminated when the BSADF sequence crosses once again the sequence of critical values from above and lies below it afterwards. If the BSADF sequence goes above the critical values once again, then this is labelled as the beginning of a new explosive episode.

It is obvious that the dating procedure erroneously identifies explosive episodes quite frequently. Even in small samples, and for data that evolve as a random walk with a weak, local-to-zero drift, explosive episodes are detected in a high proportion of the Monte Carlo replications. For the largest sample size considered, one or more explosive episodes with a

¹⁸In a related context, issues arising from the multiple application of statistical tests designed to detect explosive behavior as new observations become available are discussed in [Homm and Breitung \(2012\)](#) and [Astill, Harvey, Leybourne, Sollis, and Taylor \(2018\)](#).

¹⁹The BSADF critical values were obtained using PSY's Matlab code, which is available at <https://sites.google.com/site/shupingshi/home/codes>.

T	$k = 1$	$k = 2$	$k = 3$	$k = 4$	$k = 5$	$k = 6$	$k = 9$	$k = 12$
Estimated Probability of Detecting Explosiveness								
100	0.712	0.500	0.350	0.250	0.187	0.139	0.071	0.038
200	0.835	0.676	0.549	0.442	0.363	0.300	0.174	0.111
400	0.919	0.827	0.732	0.645	0.576	0.511	0.358	0.257
800	0.972	0.926	0.875	0.815	0.762	0.711	0.566	0.453
1600	0.990	0.971	0.951	0.929	0.904	0.875	0.784	0.687
Average Number of Explosive Episodes								
100	1.520	0.767	0.467	0.306	0.215	0.154	0.074	0.039
200	2.581	1.398	0.931	0.667	0.501	0.390	0.203	0.122
400	4.222	2.456	1.710	1.287	1.022	0.832	0.492	0.322
800	6.921	4.191	3.062	2.386	1.932	1.616	1.038	0.736
1600	11.085	6.988	5.221	4.182	3.478	2.973	2.027	1.491

Table 7: Proportion of artificial samples in which at least k consecutive BSADF statistics exceed, at least once, the relevant 0.05-level critical values, and average number of explosive episodes per sample

duration of at least 12 periods are detected in 68.4% of the replications. Simple accounting reveals that it is highly likely that at least one explosive episode will be identified by the crossing rule even when the GSADF test does not reject the unit-root hypothesis.

These results demonstrate that the convenience of using conventional fixed-level (e.g., 0.05-level) critical values for the sequence of BSADF statistics comes at a significant cost. The results also offer a partial explanation for the large number of “bubbles” that are frequently identified in empirical applications on the basis of the PWY and PSY date-stamping algorithms. A simple way of controlling the number of false detections of explosiveness by the crossing rule is to decrease the common nominal level for the individual thresholds $\{\hat{C}_\beta(r_2); r_2 \in \mathcal{R}_2\}$ from β to $\beta_m := m\beta/|\mathcal{R}_2|$, where m is a predetermined integer (with $1 \leq m < |\mathcal{R}_2|$). This ensures that, when no explosive episodes are present, the probability of $BSADF(r_0, r_2)$ exceeding $\hat{C}_\beta(r_2)$ for at least m values of $r_2 \in \mathcal{R}_2$ (albeit

not necessarily consecutive) is approximately no greater than a designated value β .^{20,21}

An alternative approach is to use a suitable bootstrap procedure to calibrate the common significance level for individual thresholds so that, in the absence of explosiveness, the probability of $BSADF(r_0, r_2)$ exceeding the corresponding critical value for k or more consecutive values of $r_2 \in \mathcal{R}_2$ (for some predetermined $1 \leq k < |\mathcal{R}_2|$), at least once during the sample period, is as close as possible to a designated desired value β (e.g., $\beta = 0.05$). Such a procedure has the obvious advantage that only potential explosive episodes with a minimum duration of k periods are taken into consideration.

For some given values of (k, β) and a sufficiently large positive integer B , the calibration procedure involves the following steps:

- (i) construct B bootstrap time series $\{Y_{b,t}^*\}$, $b = 1, 2, \dots, B$, of the same length as $\{Y_t\}$, using the wild-bootstrap scheme described in Section 4.3;
- (ii) compute B collections $\{BSADF_b^*(r_0, r_2); r_2 \in \mathcal{R}_2\}$, $b = 1, 2, \dots, B$, of BSADF values by applying the definition of $BSADF(r_0, r_2)$ to each $\{Y_{b,t}^*\}$ in place of $\{Y_t\}$;
- (iii) for each $r_2 \in \mathcal{R}_2$ and any $\lambda \in (0, 1)$, obtain the λ -level threshold for $BSADF(r_0, r_2)$ as the $\lfloor (B+1)(1-\lambda) \rfloor$ -th largest of the values $BSADF_1^*(r_0, r_2), \dots, BSADF_B^*(r_0, r_2)$;
- (iv) for a grid of values of λ , compute the proportion $\pi_k^*(\lambda)$ of bootstrap time series which exhibit at least one explosive episode at level λ , defining an explosive episode as a segment of $\{Y_{b,t}^*\}$ in which the bootstrap BSADF statistics exceed the corresponding λ -level thresholds for at least k consecutive values of r_2 ;

²⁰This is easily seen by noting that $|\mathcal{Q}_{\beta m}| = \sum_{r_2 \in \mathcal{R}_2} I\{BSADF(r_0, r_2) > \hat{C}_{\beta m}(r_2)\}$, with $I\{A\}$ denoting the indicator variable of the event A , and hence, by Markov's inequality, $P(|\mathcal{Q}_{\beta m}| \geq m) \leq m^{-1}E(|\mathcal{Q}_{\beta m}|) \approx m^{-1}\beta m |\mathcal{R}_2| = \beta$.

²¹Repeating the simulation experiments after setting $m = \lfloor 0.05(|\mathcal{R}_2|) \rfloor$ and $\beta = 0.05$, we found that the proportion of artificial time series for which the BSADF statistics exceed, at least once, their threshold values for a minimum of k consecutive periods decreases considerably, especially for $k \geq 6$. The rationale behind this adjustment is that one may be willing to accept at least 5% of the BSADF statistics exceeding their thresholds in the absence of explosiveness, provided the probability of such an outcome is controlled. For example, when $T = 400$, the proportion in question is 0.053 and 0.0067 for $k = 4$ and $k = 12$, respectively. When using 0.05-level critical values obtained via the bootstrap procedure discussed in Phillips and Shi (2020) (with $\tau_b = 24$, in their notation), the corresponding figures are 0.442 and 0.179, respectively, suggesting that the procedure is not particularly effective in controlling the probability of false detections of explosiveness. Detailed results are available upon request.

T	No Calibration		Calibration	
	Proportion	Average	Proportion	Average
100	0.017	1.046	0.020	1.087
200	0.065	1.046	0.023	1.087
400	0.227	1.189	0.036	1.056
800	0.414	1.370	0.026	1.192
1600	0.661	2.082	0.042	1.310

Table 8: Proportion of artificial samples in which at least 12 consecutive BSADF statistics exceed, at least once, the relevant wild-bootstrap critical values, and average number of explosive episodes per sample ($\beta = 0.05$)

(v) find the value $\bar{\lambda}$ for which $\pi_k^*(\bar{\lambda}) = \beta$;

(vi) use $\bar{\lambda}$ as the calibrated significance level for individual thresholds in the date-stamping algorithm.

Table 8 contains results from simulation experiments designed to shed light on the properties of the bootstrap-based calibration procedure. Specifically, we report the proportion of artificial samples for which the BSADF statistics exceed, at least once in each sample, the relevant wild-bootstrap critical values (obtained with $B = 10,000$ and $\beta = 0.05$) for a minimum of $k = 12$ consecutive values of r_2 , with and without calibration of the significance level; we also report the average number of explosive episodes per time series detected by the PSY crossing rule. As before, results are based on 1000 artificial time series generated according to (1), with $\delta = \eta = 1$ and $\{u_t\}$ being i.i.d. $N(0, 1)$ variates.

The bootstrap-based calibration procedure is quite successful at controlling the probability of erroneously detecting one or more explosive episodes using the sequence of BSADF statistics, the Monte Carlo estimates of the probability in question being somewhat smaller than the target value 0.05. A crossing rule using calibrated bootstrap critical values identifies, on average, one explosive episode in each artificial sample.

It should be pointed out that, in the preceding discussion, the dating algorithm based on the sequence of BSADF statistics has been treated primarily as a means of detect-

ing deviations from the unit-root hypothesis and of identifying the number of explosive episodes. In applications, date-stamping is often a combined procedure resulting from first implementing the GSADF test and then, if the test rejects the unit-root hypothesis, using the BSADF statistics to estimate the origination and termination dates of explosive episodes. To reflect this practice in our analysis, we revisit our earlier simulation experiments, based on (1) with $\delta = \eta = 1$, and report in Table 9 the proportion of artificial time series for which at least one explosive episode is detected, as well as the average number of explosive episodes identified by the crossing rule in each artificial time series. Here, an explosive episode is defined as the event that the 0.05-level GSADF test rejects the unit-root hypothesis and the BSADF sequence exceeds the relevant 0.05-level critical values for at least k consecutive values of r_2 ; if the GSADF test does not reject, then the number of explosive episodes is considered to be zero, regardless of the values of the BSADF sequence. To ensure that the GSADF test does not suffer from significant level distortions, we rely on calibrated wild-bootstrap critical values, obtained as in Section 4.4 (with $B = 2N = 1000$); BSADF critical values are obtained by means of the simulation-based procedure described in Section 5.1 (with $M = 2000$).

As can be seen in Table 9, when artificial time series for which the bootstrap-assisted GSADF test does not reject the unit-root hypothesis are classified as non-explosive, even if individual BSADF statistics exceed the corresponding thresholds, the estimated probability of detecting one or more explosive episodes is very close to the nominal 0.05 value and the average number of explosive episodes detected is less than one in all cases. In light of the simulation evidence discussed in Section 4.4, such results are not very surprising, and reconfirm our earlier observation that calibrated bootstrap-assisted GSADF tests tend to have finite-sample rejection probabilities under the unit-root hypothesis which are close to their nominal level.²²

A note of caution, however, is warranted. Within the subset of artificial time series for which the GSADF statistic exceeds its calibrated wild-bootstrap critical value, the

²²In contrast to the DGP used in the simulations in Section 4.4, which has a fixed non-zero drift, the DGP used here has a local-to-zero drift. It is clear, therefore, that the calibrated wild-bootstrap procedure is capable of delivering GSADF tests with minimal level distortions under vanishing-drift conditions too.

T	$k = 1$	$k = 2$	$k = 3$	$k = 4$	$k = 5$	$k = 6$	$k = 9$	$k = 12$
Estimated Probability of Detecting Explosiveness								
100	0.050	0.050	0.048	0.043	0.039	0.035	0.027	0.015
200	0.045	0.045	0.045	0.044	0.043	0.042	0.034	0.025
400	0.051	0.051	0.051	0.051	0.051	0.051	0.047	0.042
800	0.050	0.050	0.050	0.050	0.050	0.050	0.050	0.048
1600	0.048	0.048	0.048	0.048	0.048	0.048	0.048	0.048
Average Number of Explosive Episodes								
100	0.127	0.089	0.072	0.057	0.049	0.042	0.029	0.016
200	0.168	0.112	0.088	0.077	0.068	0.060	0.043	0.029
400	0.292	0.185	0.147	0.123	0.107	0.095	0.069	0.054
800	0.467	0.301	0.239	0.196	0.165	0.147	0.111	0.092
1600	0.660	0.421	0.322	0.272	0.236	0.212	0.160	0.131

Table 9: Proportion of artificial samples in which the 0.05-level GSADF test rejects and at least k consecutive BSADF statistics exceed, at least once, the relevant 0.05-level critical values, and average number of explosive episodes detected per sample

estimated probability of one or more explosive episodes being erroneously detected by the BSADF-based crossing rule remains high, as does the average number of such episodes identified in each time series, especially in large samples – the relevant figures (not reported here) are, in fact, somewhat larger than those in Table 7. Put differently, implementing the date-stamping procedure only if the unit-root hypothesis is rejected on the basis of the GSADF test, does not imply that the expected proportion of originating explosive episodes that are erroneously identified as such by the crossing rule will be close to the nominal level of individual BSADF tests. The bootstrap-based calibration procedure discussed earlier in this section provides a practical way of controlling such errors.

In concluding this section, we note that, as the highlighted issue of erroneous discoveries of explosiveness is associated with the way critical values for BSADF statistics are used in the crossing principle, it is unrelated to the separate issue of possible misspecification of the drift characteristics of the data, which was discussed in Sections 3 and 4.

6 Empirical analysis

In this section, we investigate whether explosive episodes are present in U.S. stock prices using Robert Shiller’s data set of monthly real prices and real dividends associated with the S&P 500 stock index.²³ Our analysis focuses on logarithmic prices (as in PWY and Phillips, Shi, and Yu (2014)) and on the price–dividend ratio (as in PSY) for the period from March 1927 to June 2020 (a total of 1120 data points). Logarithmic prices may perhaps appear to be less attractive than the price–dividend ratio when examining asset-price explosiveness since present-value and bubble solutions to equations like (8) or (13) are typically associated with raw prices.²⁴ However, following Campbell and Shiller (1989), log-linear approximations of these equations are commonly used in theoretical and applied work, notwithstanding the fact that the use of such approximations (with present-value and bubble components expressed in logarithms, as, e.g., in PWY) requires careful consideration in the presence of explosive behavior. Since a discussion of the appropriateness or otherwise of log-linear approximations is beyond the scope of this paper, logarithmic prices ($\ln P_t$), logarithmic dividends ($\ln D_t$), and the price–dividend ratio (P_t/D_t) are considered in the sequel.

PSY argue against using a large value for ℓ in ADF regression equations such as (2) because this typically results in SADF and GSADF tests having actual rejection probabilities under the unit-root hypothesis that are much larger than the nominal level. Following their recommendation, we use a first-order autoregressive model ($\ell = 0$) in the implementation of the tests (the model also used in their empirical application).

The values of the SADF and GSADF statistics (with $r_0 = 0.01 + 1.8T^{-1/2} = 0.06378$) for the three time series under consideration are shown in Table 10, along with finite-sample critical values for right-tailed tests associated with $\eta = 1$ and asymptotic critical values associated with $\eta = 0$ (for three significance levels). We also report bootstrap crit-

²³The data is available at www.econ.yale.edu/shiller/data.htm.

²⁴By contrast, logarithmic prices arise naturally in analyses of hyperinflations or foreign-exchange rates, monetary models of which have similar structure to (8) and (13), with P_t corresponding to the logarithm of the aggregate price level or the logarithm of the nominal exchange rate and D_t corresponding to the logarithm of nominal money stock or the logarithm of a market-fundamentals variable that involves relative money supply and real income.

	SADF			GSADF		
$\ln P_t$	0.3456			2.9949		
$\ln D_t$	2.6474			7.8558		
P_t/D_t	3.1271			4.1603		
Critical Values	0.10	0.05	0.01	0.10	0.05	0.01
$\eta = 1$	1.2896	1.5957	1.9859	2.1900	2.4100	2.8700
$\eta = 0$	2.3500	2.7000	3.3600	3.9900	4.2300	4.6000
Bootstrap – $\ln P_t$	1.2243	1.5649	2.1673	3.1662	3.5054	4.3846
Bootstrap – $\ln D_t$	2.2967	2.7958	3.9629	4.2177	4.6505	5.6782
Bootstrap – P_t/D_t	2.0947	2.6113	3.7806	3.8694	4.4734	5.6869
Calibrated – $\ln P_t$	1.1904	1.4597	1.9667	3.0488	3.2706	4.1202
	(0.1118)	(0.0635)	(0.0200)	(0.1159)	(0.0775)	(0.0155)
Calibrated – $\ln D_t$	2.3749	2.9619	4.1361	4.3651	4.9998	6.2308
	(0.0886)	(0.0408)	(0.0068)	(0.0777)	(0.0293)	(0.0041)
Calibrated – P_t/D_t	1.7249	2.3638	4.0741	3.8042	4.5757	5.8166
	(0.1326)	(0.0638)	(0.0072)	(0.1070)	(0.0434)	(0.0083)

Table 10: SADF and GSADF statistics for logarithmic real stock prices ($\ln P_t$), logarithmic real dividends ($\ln D_t$), and the real price–dividend ratio (P_t/D_t), and critical values for right-tailed unit-root tests. Figures in parentheses are calibrated significance levels ($\tilde{\lambda}$).

ical values obtained from $B = 1000$ wild-bootstrap replicates of the SADF and GSADF statistics, respectively (see Section 4.3); additionally, calibrated bootstrap critical values are reported based on $B = N = 1000$ first-round and second-round wild-bootstrap replicates of the two test statistics (see Section 4.4). Recall that our wild-bootstrap procedure does not restrict the drift (for either the observed or the bootstrap data) to be zero, or local-to-zero, under the null hypothesis.

As can be seen in Table 10, the differences between bootstrap critical values for the GSADF test (with or without calibration) and critical values simulated under the assumption of a local-to-zero drift ($\eta = 1$) are quite substantial. The unit-root hypothesis is rejected for logarithmic prices when using the latter set of critical values, but is not rejected using the former (or when using asymptotic critical values associated with $\eta = 0$); the unit-root hypothesis cannot be rejected for $\ln P_t$ on the basis of the SADF test regardless of what critical values are used. In the case of the price–dividend ratio, the null

hypothesis is rejected, at the 10% level, using either of the two tests and any of the four different critical values; referring the SADF statistic to 0.05-level critical values also leads to rejection (but only the critical value associated with $\eta = 1$ leads to rejection by the GSADF test at the 5% level). Lastly, the unit-root hypothesis is rejected, at the 1% level, for logarithmic dividends on the basis of the GSADF test; the SADF test also rejects, at the 10% level, using any of the reported critical values (with only the critical value associated with $\eta = 1$ leading to rejection at the 5% level).

As we have argued, such differences are the result of the maintained assumptions about the drift characteristics of the data. Under the DGP in (1) with $\delta = \eta = 1$, the implicit drift used to simulate critical values for unit-root tests is $1/1120 \approx 0.00089$. However, the estimated drift in logarithmic prices under the null hypothesis of a unit root is 0.0025, about three times higher. In view of our earlier simulation findings, it is likely, therefore, that the use of critical values from a localized-drift DGP would lead to rejection of the unit-root hypothesis.

Turning attention to the date-stamping of explosive episodes, it is important to deal with the difficulties that arise from the use of the sequential BSADF statistics. In order to do so, we use the bootstrap-based procedure described in Section 5.2 to calibrate the common significance level for individual thresholds (critical values) so that, in the absence of explosiveness, and irrespective of the outcome of the GSADF test, the (estimated) probability of at least 12 consecutive BSADF statistics exceeding their corresponding thresholds, at least once during the sample period, is as close as possible to 0.05.

Figure 1 shows the sequence of BSADF statistics and the corresponding 0.05-level and $\bar{\lambda}$ -level wild-bootstrap critical values (obtained from $B = 10,000$ bootstrap replications). The plot highlights the potential difficulty discussed in Section 5, namely that, in a given sample, one may fail to reject the unit-root hypothesis on the basis of the GSADF test while at the same time several apparent explosive episodes may be identified using a crossing rule based on the collection of BSADF statistics with 5% nominal significance level. This is precisely what is observed here: even though the GSADF test fails to reject the unit-root hypothesis for logarithmic prices, the crossing principle finds evidence of at




Figure 1: Sequence of BSADF statistics (blue), 0.05-level bootstrap critical values (red, dotted), calibrated bootstrap critical values (red, dashed), and logarithmic prices (black, right axis).




Figure 2: The top panel shows the time series corresponding to the price–dividend ratio (black, right axis), the sequence of BSADF statistics (blue), 0.05-level bootstrap critical values (red, dotted), and calibrated bootstrap critical values (red, dashed). The bottom panel shows the same information for logarithmic dividends.

least two explosive episodes in the data (on the basis of 0.05-level critical values), one during the 1950s and the other during the late 1990s.²⁵ A similar pattern is observed in the top panel of Figure 2 for the price–dividend ratio. As noted earlier, this issue arises because of the way critical values for the BSADF statistics are used for dating purposes and is independent of whether one relies on critical values obtained by means of the Monte Carlo algorithm described in Section 5.1 or by a bootstrap procedure (as is the case here).

A rather different picture emerges when calibrated $\bar{\lambda}$ -level critical values are used for the BSADF statistics. Specifically, only three crossings are detected for logarithmic prices, none of which lasts for more than five months.²⁶ The outcome of the dating algorithm for the price–dividend ratio is qualitatively similar.²⁷ The calibration-based procedure evidently reduces the number of episodes identified as explosive, when such episodes are defined as periods in which the BSADF statistics exceed the corresponding thresholds for at least 12 consecutive months. In fact, under this definition, no explosive episodes are detected in either logarithmic prices or the price–dividend ratio. It is important to note that defining what an explosive episode is more liberally (for example, by requiring that it lasts 6 months or more) would not necessarily imply that more explosive episodes would be identified. The reason for this is that the calibrated significance level $\bar{\lambda}$ would adjust to ensure that, in the absence of explosiveness, the BSADF statistics exceed the corresponding thresholds for the required minimum number of consecutive periods in only 5% (approximately) of the bootstrap samples.²⁸

Somewhat counter-intuitively, as can be seen in Table 10 and in the lower panel of

²⁵Driffill and Sola (1998) attribute the first episode to stochastic regime-switching in the DGP of real logarithmic dividends rather than a bubble.

²⁶The periods in which the BSADF statistics go above the $\bar{\lambda}$ -level critical values are 1995:7–1995:9, 1998:3–1998:7, and one lasting only one month, 1955:4. As none of these lasts for 12 months or more, no explosive episodes are identified under the definition used here ($k = 12$).

²⁷The periods in which the BSADF statistics go above the $\bar{\lambda}$ -level critical values are 1954:12–1955:4, 1955:6–1955:7, 1955:9, 1987:8, 1998:2–1998:7, and 2000:8, none of them extending for more than five months.

²⁸We note that SADF and GSADF results for logarithmic prices and the price–dividend ratio are not substantially different when second-order autoregressive models ($\ell = 1$) are used in the implementation of the tests, and lead to the same overall conclusions. Similarly, BSADF tests with bootstrap-calibrated significance levels identify no explosive episodes lasting for at least a year. Detailed results are available upon request.

Figure 2, the only time series in which clear evidence of multiple periods of explosiveness is found is that of real dividends. Of course, since $\ln P_t = \ln(P_t/D_t) + \ln D_t$, explosiveness in either $\ln D_t$ or $\ln(P_t/D_t)$ would be inherited by $\ln P_t$. However, it is rather difficult to justify explosiveness of logarithmic dividends from a theoretical point of view. In much of the related theoretical and empirical work, logarithmic dividends are considered to be a martingale with a drift component which is fixed, slowly varying, or subject to discrete shifts (see, e.g., [Froot and Obstfeld \(1991\)](#), [Driffill and Sola \(1998\)](#), and [Pettenuzzo, Sabbatucci, and Timmermann \(2020\)](#)). One possible explanation for the presence of multiple explosive episodes in logarithmic dividends may lie with the nature of the dividend data. The data used here is monthly, but as Robert Shiller points out on his website, monthly data are obtained from quarterly data via linear interpolation. This can potentially introduce non-trivial dynamics into the dividend series.²⁹ Furthermore, interpolation tends to reduce the estimated variance of innovations in regression equations, thus resulting in studentized statistics that are biased towards more extreme values. These problems may also affect results for the price–dividend ratio since it is constructed using monthly dividends.

7 Conclusion

This paper has investigated two important issues that arise in the practical implementation of the widely used PWY and PSY procedures for detecting explosive behavior in segments of a time series using recursive right-tailed unit-root tests.

The first issue relates to the assumption of a local-to-zero drift under the unit-root hypothesis, which typically underlies the implementation of the PWY and PSY procedures. Referring SADF and GSADF statistics to critical values that are obtained under the assumption of a weak and asymptotically vanishing drift can lead to over-rejection of the unit-root hypothesis in favor of an explosive alternative when the drift in the data is not negligible relative to the martingale component. Our simulation experiments have revealed that, for sample sizes and parameter values that match the properties of time series such as stock prices, over-rejection of the unit-root hypothesis can be quite severe. Allowing for

²⁹As not all firms with common stocks report dividends in each quarter, a quarter-on-quarter comparison is not free of problems (see [Pettenuzzo, Sabbatucci, and Timmermann \(2020\)](#)).

a non-vanishing dominant drift under the null hypothesis is not without difficulties either since, unless the drift is substantial, recursive right-tailed unit-root tests tend to under-reject. The use of a suitable bootstrap procedure to obtain critical values for the tests is advisable and does not add much in terms of computational complexity compared to the PSY simulation procedure that is typically used to compute finite-sample critical values. However, even this resampling procedure can be only partially successful at controlling the tendency for under-rejection of the unit-root hypothesis in the presence of a non-vanishing drift, especially so in the case of GSADF tests. We have discussed how such difficulties may be overcome by using a bootstrap-based calibration procedure to control errors in the rejection probabilities of the tests.

The second issue relates to the dating of the origination and termination dates of explosive episodes based on the PSY crossing rule involving the sequence of BSADF statistics. We have argued that the standard practice of using a conventional level of significance (e.g., 5%) for the critical values (thresholds) associated with each BSADF statistic does not guarantee control of the probability of one or more explosive episodes being erroneously identified as such. Our simulations have revealed that the probability of detecting at least one such episode using the crossing principle, when no explosive episodes are present, can be substantially larger than the nominal level associated with the BSADF threshold values. Although requiring explosive episodes to have a suitable minimum duration reduces the probability of false discoveries of explosiveness, the dating procedure still erroneously identifies the beginning and end of explosive episodes with high probability. Such findings, together with those highlighted in the previous paragraph, offer a possible explanation for the relatively large number of “bubbles” in real-world time series that are frequently reported in empirical applications that make use of the PWY and PSY inferential procedures. These difficulties must be taken into account in applied work, in which it is advisable to use suitable adjustments for the significance level of individual BSADF thresholds to control the probability of these thresholds being (wrongly) crossed only an acceptable number of times. We have considered such adjustments based on a bootstrap-assisted calibration technique.

In our analysis of monthly U.S. real stock prices, we have found that the GSADF test rejects the null hypothesis of a unit root in favor of an explosive alternative when critical values obtained under the assumption of a local-to-zero drift are used, but does not reject when using critical values obtained by means of a wild-bootstrap procedure which allows for a non-vanishing drift. Despite the latter result, using (bootstrap) critical values for BSADF statistics to date-stamp explosive episodes, we identify at least two such episodes lasting a year or more, one during the 1950s and the other during the late 1990s. No explosive episodes are found when the significance level of the critical values used in the dating procedure is suitably calibrated to control the probability of erroneous detections of explosiveness.

In closing, it should be stressed that recursive supremum-ADF procedures are undoubtedly valuable techniques for identifying explosive bubble-like behavior and have deservedly attracted much attention in applied research. Nevertheless, they must be used with great care as results can vary considerably depending on the way the procedures are implemented and on the maintained assumptions about the underlying data-generating mechanism.

A Appendix

A.1 Price–dividend ratio

To obtain the market-fundamentals solution to (8) under (6)–(7), we need to evaluate $E_t(D_{t+j})$ for $j \geq 1$. To this end, putting $d_t := \ln D_t$ and iterating (6) and (7) backwards, we have

$$d_{t+j} = d_t + \sum_{i=1}^j \mu_{t+i} + \sigma_d \sum_{i=1}^j \varepsilon_{t+i},$$

with

$$\mu_{t+i} = \mu(1 - \phi^i) + \phi^i \mu_t + \sigma_\mu \sum_{h=0}^{i-1} \phi^h \zeta_{t+i-h},$$

and hence

$$d_{t+j} = d_t + \sum_{i=1}^j \left(\mu(1 - \phi^i) + \phi^i \mu_t + \sigma_\mu \sum_{h=0}^{i-1} \phi^h \zeta_{t+i-h} \right) + \sigma_d \sum_{i=1}^j \varepsilon_{t+i}.$$

Therefore, d_{t+j} is conditionally Gaussian, given $\{d_n; n \leq t\}$, with

$$\begin{aligned}\mathbb{E}_t(d_{t+j}) &= d_t + \sum_{i=1}^j [\mu(1 - \phi^i) + \phi^i \mu_t] = d_t + j\mu + (\mu_t - \mu) \sum_{i=1}^j \phi^i \\ &= d_t + j\mu + \frac{\phi(1 - \phi^j)(\mu_t - \mu)}{1 - \phi},\end{aligned}$$

and

$$\begin{aligned}\mathbb{E}_t[\{d_{t+j} - \mathbb{E}_t(d_{t+j})\}^2] &= \sigma_\mu^2 \sum_{i=1}^j \sum_{h=0}^{i-1} \phi^{2h} + j\sigma_d^2 = \sigma_\mu^2 \sum_{i=1}^j \left(\frac{1 - \phi^{2i}}{1 - \phi^2} \right) + j\sigma_d^2 \\ &= \frac{\sigma_\mu^2}{1 - \phi^2} \left(j - \frac{\phi^2(1 - \phi^{2j})}{1 - \phi^2} \right) + j\sigma_d^2.\end{aligned}$$

Consequently,

$$\begin{aligned}\mathbb{E}_t(D_{t+j}) &= \exp \left(d_t + j\mu + \frac{\phi(1 - \phi^j)(\mu_t - \mu)}{1 - \phi} + \frac{\sigma_\mu^2}{2(1 - \phi^2)} \left\{ j - \frac{\phi^2(1 - \phi^{2j})}{1 - \phi^2} \right\} + \frac{j\sigma_d^2}{2} \right) \\ &= D_t \exp \left(j \left[\mu + \frac{\sigma_\mu^2}{2(1 - \phi^2)} + \frac{\sigma_d^2}{2} \right] + \frac{\phi(1 - \phi^j)(\mu_t - \mu)}{1 - \phi} - \frac{\phi^2(1 - \phi^{2j})\sigma_\mu^2}{2(1 - \phi^2)^2} \right),\end{aligned}$$

which, together with $P_t = \sum_{j=1}^{\infty} e^{-\rho j} \mathbb{E}_t(D_{t+j})$, yields the result in (9).

A.2 Parameter calibration

The model for dividend growth of [Pettenuzzo, Sabbatucci, and Timmermann \(2020\)](#), adapted to monthly frequency (also used in PWY and PSY) and excluding jump and transitory components, is given by

$$\ln(D_t/D_{t-12}) = \tilde{\mu}_t + \tilde{\sigma}_d \zeta_{d,t}, \quad (\text{A.1})$$

$$\tilde{\mu}_t - \tilde{\mu} = \tilde{\phi}(\tilde{\mu}_{t-1} - \tilde{\mu}) + \tilde{\sigma}_\mu \zeta_{\mu,t}, \quad (\text{A.2})$$

where $\{\zeta_{d,t}\}$ and $\{\zeta_{\mu,t}\}$ are independent sequences of i.i.d. $N(0, 1)$ random variables. This is a model for the one-year growth rate instead of the monthly growth rate. To explain how to use the model for calibration, suppose monthly logarithmic dividends satisfy (6)–(7), but we are able to estimate only the parameters $(\tilde{\sigma}_d, \tilde{\mu}, \tilde{\phi}, \tilde{\sigma}_\mu)$ in (A.1)–(A.2); estimates $(\tilde{\sigma}_d^*, \tilde{\mu}^*, \tilde{\phi}^*, \tilde{\sigma}_\mu^*)$ of these parameters may be obtained by the Bayesian approach of [Pettenuzzo, Sabbatucci, and Timmermann \(2020\)](#) or by classical maximum-likelihood methods that rely on Kalman filtering.

Since

$$\ln(D_t/D_{t-12}) = \sum_{j=0}^{11} \mu_{t-j} + \sigma_d \sum_{j=0}^{11} \varepsilon_t,$$

with $E(\sum_{j=0}^{11} \mu_{t-j}) = 12\mu$ and $\text{Var}(\sigma_d \sum_{j=0}^{11} \varepsilon_{t-j}) = 12\sigma_d^2$, the parameters μ and σ_d may be estimated as $\tilde{\mu}^*/12$ and $\tilde{\sigma}_d^*/\sqrt{12}$, respectively. Next, note that the autocovariance matrix $\mathbf{\Gamma} := \{\text{Cov}(\mu_t, \mu_{t-|i-j|})\}_{i,j=1}^{13}$ is the Toeplitz matrix $\mathbf{\Gamma} = \sigma_\mu^2(1 - \phi^2)^{-1}\mathbf{\Phi}$, where

$$\mathbf{\Phi} := \begin{pmatrix} 1 & \phi & \phi^2 & \dots & \phi^{12} \\ \phi & 1 & \phi & \dots & \phi^{11} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \phi^{12} & \phi^{11} & \phi^{10} & \dots & 1 \end{pmatrix}.$$

Hence,

$$\tilde{\phi} = \frac{\text{Cov}(\tilde{\mu}_t, \tilde{\mu}_{t-1})}{\text{Var}(\tilde{\mu}_t)} = \frac{(\mathbf{1}', 0)\mathbf{\Phi}(0, \mathbf{1}')'}{(\mathbf{1}', 0)\mathbf{\Phi}(\mathbf{1}', 0)'}, \quad (\text{A.3})$$

and

$$\begin{aligned} \tilde{\sigma}_\mu^2 &= \text{Var}(\tilde{\mu}_t) - \tilde{\phi}\text{Cov}(\tilde{\mu}_t, \tilde{\mu}_{t-1}) \\ &= \left(\frac{\sigma_\mu^2}{1 - \phi^2} \right) \left\{ (\mathbf{1}', 0)\mathbf{\Phi}(\mathbf{1}', 0)' - \frac{[(\mathbf{1}', 0)\mathbf{\Phi}(0, \mathbf{1}')']^2}{(\mathbf{1}', 0)\mathbf{\Phi}(\mathbf{1}', 0)'} \right\}, \end{aligned} \quad (\text{A.4})$$

where $\mathbf{1}$ is a 12-dimensional column vector of all ones. Equations (A.3) and (A.4) may be solved for (ϕ, σ_μ) to obtain estimates of these parameters based on $(\tilde{\phi}^*, \tilde{\sigma}_\mu^*)$.

Finally, the discount rate ρ is calibrated using the monthly real total returns series in Robert Shiller's database. Specifically, ρ is set to match the average real total monthly return during the period 1973–2016, which is the same as the post-1973 period considered in [Pettenuzzo, Sabbatucci, and Timmermann \(2020\)](#).

A.3 Asymptotics under a non-vanishing drift

In this appendix, we obtain the asymptotic distribution of the GSADF statistic (4) and of its wild-bootstrap analog when there is a fixed, non-zero drift under the null hypothesis, and establish the asymptotic correctness of the bootstrap-assisted GSADF test. Throughout the appendix, the arrows \rightsquigarrow and \rightarrow_p are used to denote convergence in distribution and convergence in probability, respectively; $O_p(1)$ and $o_p(1)$ signify being bounded in probability and converging in probability to zero, respectively, under the probability measure

\mathbf{P} governing the sample $\mathcal{Y} := \{Y_t; t = 0, 1, \dots, T\}$; \mathbf{P}^* and \mathbf{E}^* denote bootstrap probability and expectation, respectively, conditional on \mathcal{Y} ; for a sequence of bootstrap statistics $\{V_T^*\}$, $V_T^* = O_p^*(1)$ means that $\mathbf{P}^*(|V_T^*| > l_T) = o_p(1)$ for any constants $\{l_T\}$ tending to infinity, and $V_T^* = o_p^*(1)$ means that $\mathbf{P}^*(|V_T^*| > \epsilon) = o_p(1)$ for any $\epsilon > 0$; $V_T^* \rightsquigarrow_p V$ means that the distance between the conditional distribution of V_T^* , given \mathcal{Y} , and the distribution of V is $o_p(1)$, for any distance metrizing convergence in distribution to V (or, equivalently, that this distance converges almost surely to zero along some subsequence of any arbitrary subsequence of natural numbers). Unless otherwise stated, all convergence and order relations are to be understood for T tending to infinity.

For the sake of simplicity, the DGP is assumed to be

$$\Delta Y_t = \alpha + u_t, \quad \alpha \neq 0, \quad (\text{A.5})$$

for $t = 1, 2, \dots, T$, with Y_0 having a given distribution (not depending on T). It is further assumed that $\{u_t\}$ is a sequence of martingale differences (relative to some filtration) such that $\mathbf{E}(u_t^2) = \sigma^2 > 0$ for all t , $T^{-1} \sum_{t=1}^T u_t^2 \rightarrow_p \sigma^2$, and $\sup_t \mathbf{E}(u_t^4) \leq M_u < \infty$. Note that the sequence $\{u_t\}$ may be dependent (but uncorrelated) and conditionally heteroskedastic (but covariance-stationary).³⁰ The case where $\{u_t\}$ is a covariance-stationary process that admits an autoregressive representation can be handled in an analogous manner using the ADF regression (2) with $\ell \geq 1$ (without any change in the limit of the distribution of the GSADF statistic).

The asymptotic null distribution of $GSADF(r_0)$ is given in the following proposition.

Proposition A.1. *Under the DGP (A.5), $GSADF(r_0) \rightsquigarrow \Lambda$, where*

$$\Lambda := \sup_{\substack{r_1 \in [0, r_2 - r_0] \\ r_2 \in [r_0, 1]}} \left\{ \frac{\sqrt{3}}{(r_2 - r_1)^{3/2}} \left[(r_2 - r_1) \{W(r_2) + W(r_1)\} - 2 \int_{r_1}^{r_2} W(s) ds \right] \right\},$$

and $\{W(s)\}$ is a standard Brownian motion on $[0, 1]$.

Proof. For some $0 < r_1 < r_2 \leq 1$, let

$$\Delta Y_t = \hat{\alpha} + \hat{\theta} Y_{t-1} + \hat{u}_t, \quad t = T_1, T_1 + 1, \dots, T_2, \quad (\text{A.6})$$

³⁰For instance, processes that belong to the class of infinite-order autoregressive conditionally heteroskedastic models satisfy these assumptions under mild conditions, as do stochastic volatility processes.

be the fitted least-squares Dickey–Fuller regression equation over observations $[Tr_1] =: T_1, \dots, [Tr_2] =: T_2$, so that

$$\begin{pmatrix} \hat{\alpha} - \alpha \\ \hat{\theta} \end{pmatrix} = \begin{pmatrix} T_2 - T_1 + 1 & \sum_{t=T_1}^{T_2} Y_{t-1} \\ \sum_{t=T_1}^{T_2} Y_{t-1} & \sum_{t=T_1}^{T_2} Y_{t-1}^2 \end{pmatrix}^{-1} \begin{pmatrix} \sum_{t=T_1}^{T_2} u_t \\ \sum_{t=T_1}^{T_2} Y_{t-1} u_t \end{pmatrix}. \quad (\text{A.7})$$

Using standard arguments and the functional central limit theorem for martingale differences (e.g., Aldous (1978, Theorem 3)), it is straightforward to show that

$$\begin{pmatrix} T^{-1/2} \sum_{t=T_1}^{T_2} u_t \\ T^{-3/2} \sum_{t=T_1}^{T_2} Y_{t-1} u_t \end{pmatrix} \rightsquigarrow \begin{pmatrix} \sigma\{W(r_2) - W(r_1)\} \\ \alpha\sigma\{r_2 W(r_2) - r_1 W(r_1) - \int_{r_1}^{r_2} W(s) ds\} \end{pmatrix}. \quad (\text{A.8})$$

Moreover, letting $\xi_t := \sum_{j=1}^t u_j$, we have $T^{-3/2} \sum_{t=T_1}^{T_2} \xi_{t-1} = O_p(1)$, $T^{-2} \sum_{t=T_1}^{T_2} \xi_{t-1}^2 = O_p(1)$, and $T^{-5/2} \sum_{t=T_1}^{T_2} (t-1)\xi_{t-1} = O_p(1)$. Consequently,

$$\begin{aligned} T^{-2} \sum_{t=T_1}^{T_2} Y_{t-1} &= T^{-2} \sum_{t=T_1}^{T_2} \{Y_0 + \alpha(t-1) + \xi_{t-1}\} = \alpha T^{-2} \sum_{t=T_1}^{T_2} (t-1) + o_p(1) \\ &\rightarrow_p \alpha \int_{r_1}^{r_2} s ds = \frac{1}{2} \alpha (r_2^2 - r_1^2), \end{aligned} \quad (\text{A.9})$$

and

$$\begin{aligned} T^{-3} \sum_{t=T_1}^{T_2} Y_{t-1}^2 &= T^{-3} \sum_{t=T_1}^{T_2} \{Y_0 + \alpha(t-1) + \xi_{t-1}\}^2 = \alpha^2 T^{-3} \sum_{t=T_1}^{T_2} (t-1)^2 + o_p(1) \\ &\rightarrow_p \alpha^2 \int_{r_1}^{r_2} s^2 ds = \frac{1}{3} \alpha^2 (r_2^3 - r_1^3). \end{aligned} \quad (\text{A.10})$$

Hence, using the scaling matrix $\mathbf{Y}_T := \text{diag}(T^{1/2}, T^{3/2})$, (A.7) becomes

$$\mathbf{Y}_T \begin{pmatrix} \hat{\alpha} - \alpha \\ \hat{\theta} \end{pmatrix} = \left[\mathbf{Y}_T^{-1} \begin{pmatrix} T_2 - T_1 + 1 & \sum_{t=T_1}^{T_2} Y_{t-1} \\ \sum_{t=T_1}^{T_2} Y_{t-1} & \sum_{t=T_1}^{T_2} Y_{t-1}^2 \end{pmatrix} \mathbf{Y}_T^{-1} \right]^{-1} \mathbf{Y}_T^{-1} \begin{pmatrix} \sum_{t=T_1}^{T_2} u_t \\ \sum_{t=T_1}^{T_2} u_t Y_{t-1} \end{pmatrix},$$

or, equivalently,

$$\begin{pmatrix} T^{1/2}(\hat{\alpha} - \alpha) \\ T^{3/2}\hat{\theta} \end{pmatrix} = \begin{pmatrix} T^{-1}(T_2 - T_1 + 1) & T^{-2} \sum_{t=T_1}^{T_2} Y_{t-1} \\ T^{-2} \sum_{t=T_1}^{T_2} Y_{t-1} & T^{-3} \sum_{t=T_1}^{T_2} Y_{t-1}^2 \end{pmatrix}^{-1} \begin{pmatrix} T^{-1/2} \sum_{t=T_1}^{T_2} u_t \\ T^{-3/2} \sum_{t=T_1}^{T_2} u_t Y_{t-1} \end{pmatrix}. \quad (\text{A.11})$$

In view of (A.8)–(A.10), an application of Slutsky’s theorem to (A.11) yields

$$\begin{pmatrix} T^{1/2}(\hat{\alpha} - \alpha) \\ T^{3/2}\hat{\theta} \end{pmatrix} \rightsquigarrow \mathbf{A}^{-1} \begin{pmatrix} \sigma\{W(r_2) - W(r_1)\} \\ \alpha\sigma\{r_2 W(r_2) - r_1 W(r_1) - \int_{r_1}^{r_2} W(s) ds\} \end{pmatrix},$$

where

$$\mathbf{A} := \begin{pmatrix} r_2 - r_1 & \frac{1}{2} \alpha (r_2^2 - r_1^2) \\ \frac{1}{2} \alpha (r_2^2 - r_1^2) & \frac{1}{3} \alpha^2 (r_2^3 - r_1^3) \end{pmatrix}.$$

Hence, $T^{3/2}\hat{\theta} \rightsquigarrow Z$, where

$$\begin{aligned} Z &:= \frac{1}{\det \mathbf{A}} \left[-\frac{1}{2}\alpha\sigma(r_2^2 - r_1^2)\{W(r_2) - W(r_1)\} + \alpha\sigma(r_2 - r_1)\{r_2W(r_2) - r_1W(r_1) - \int_{r_1}^{r_2} W(s)ds\} \right] \\ &= \frac{12\sigma}{\alpha(r_2 - r_1)^3} \left[-\frac{1}{2}(r_2 + r_1)\{W(r_2) - W(r_1)\} + r_2W(r_2) - r_1W(r_1) - \int_{r_1}^{r_2} W(s)ds \right] \\ &= \frac{12\sigma}{\alpha(r_2 - r_1)^3} \left[\frac{1}{2}(r_2 - r_1)\{W(r_2) + W(r_1)\} - \int_{r_1}^{r_2} W(s)ds \right]. \end{aligned}$$

Next, note that $ADF(r_1, r_2) = (T^{3/2}\hat{\theta})(T^{-3/2}\hat{\tau})/\hat{\sigma}$, where

$$\hat{\tau}^2 := \sum_{t=T_1}^{T_2} Y_{t-1}^2 - (T_2 - T_1 + 1)^{-1} \left(\sum_{t=T_1}^{T_2} Y_{t-1} \right)^2,$$

and $\hat{\sigma}^2 := (T_2 - T_1 - 1)^{-1} \sum_{t=T_1}^{T_2} \hat{u}_t^2$. Since it may easily be verified that $\hat{\sigma}^2 \rightarrow_p \sigma^2$ and

$$T^{-3}\hat{\tau}^2 \rightarrow_p \frac{1}{3}\alpha^2(r_2^3 - r_1^3) - \frac{1}{4}\alpha^2(r_2^2 - r_1^2)^2(r_2 - r_1)^{-1} = \frac{1}{12}\alpha^2(r_2 - r_1)^3,$$

it follows, by Slutsky's theorem, that

$$ADF(r_1, r_2) \rightsquigarrow \frac{\alpha(r_2 - r_1)^{3/2} Z}{\sqrt{12}\sigma} = \frac{\sqrt{3}}{(r_2 - r_1)^{3/2}} \left[(r_2 - r_1)\{W(r_2) + W(r_1)\} - 2 \int_{r_1}^{r_2} W(s)ds \right].$$

The last result and an argument analogous to that used in the proof of Theorem 1 of PSY allow us to conclude, by appealing to the continuous mapping theorem, that $GSADF(r_0) \rightsquigarrow \Lambda$. \square

Next, we consider the large-sample behavior of the wild-bootstrap analog $GSADF^*(r_0)$ of the $GSADF(r_0)$ statistic, defined in the same way as the latter but using $\{Y_t^*\}$, generated according to (15) (with $\ell = 0$), in place of $\{Y_t\}$. The conditional distribution of $GSADF^*(r_0)$, given \mathcal{Y} , constitutes the bootstrap estimator of the null distribution of $GSADF(r_0)$. To establish consistency of this estimator, it is enough to show that, under (A.5), the conditional distribution of $GSADF^*(r_0)$ converges weakly (in probability) to the same limit as the unconditional distribution of $GSADF(r_0)$. The next proposition shows this to be true. Recall that $\{\nu_t^*\}$ are i.i.d., independent of \mathcal{Y} , with $\mathbf{E}^*(\nu_t^*) = 0$ and $\mathbf{E}^*(\nu_t^{*2}) = 1$. It is additionally assumed that $\{\nu_t^*\}$ are chosen so that $\mathbf{E}^*(\nu_t^{*4}) \leq M_\nu < \infty$.

Proposition A.2. *Under the DGP (A.5), $GSADF^*(r_0) \rightsquigarrow_p \Lambda$.*

Proof. The bootstrap analog of the fitted regression equation (A.6) is

$$\Delta Y_t^* = \hat{\alpha}^* + \hat{\theta}^* Y_{t-1}^* + \hat{u}_t^*, \quad t = T_1, \dots, T_2,$$

with

$$\begin{pmatrix} T^{1/2}(\hat{\alpha}^* - \tilde{\alpha}) \\ T^{3/2}\hat{\theta}^* \end{pmatrix} = \begin{pmatrix} T^{-1}(T_2 - T_1 + 1) & T^{-2} \sum_{t=T_1}^{T_2} Y_{t-1}^* \\ T^{-2} \sum_{t=T_1}^{T_2} Y_{t-1}^* & T^{-3} \sum_{t=T_1}^{T_2} Y_{t-1}^{*2} \end{pmatrix}^{-1} \begin{pmatrix} T^{-1/2} \sum_{t=T_1}^{T_2} u_t^* \\ T^{-3/2} \sum_{t=T_1}^{T_2} Y_{t-1}^* u_t^* \end{pmatrix}, \quad (\text{A.12})$$

and $u_t^* := \nu_t^* \tilde{\varepsilon}_t = \nu_t^* [u_t - (\tilde{\alpha} - \alpha)]$.

Letting $J_T^* := \{\sum_{t=1}^T \mathbf{E}^*(u_t^{*2})\}^{1/2}$ and noting that $\{u_t^*\}$ are independent, conditionally on \mathcal{Y} , with $\mathbf{E}^*(u_t^*) = 0$ and $\mathbf{E}^*(u_t^{*2}) = \tilde{\varepsilon}_t^2$, we have

$$\mathbf{E}^* \left[\left(T^{-1/2} \sum_{t=1}^T u_t^* \right)^2 \right] = T^{-1} J_T^{*2} = T^{-1} \sum_{t=1}^T \tilde{\varepsilon}_t^2 = T^{-1} \sum_{t=1}^T u_t^2 + o_p(1) \rightarrow_p \sigma^2,$$

since $T^{1/2}(\tilde{\alpha} - \alpha) = O_p(1)$. Moreover, $\{u_t^*\}$ satisfy the conditional Lyapunov condition

$$(J_T^*)^{-2\kappa} \sum_{t=1}^T \mathbf{E}^*(|u_t^*|^{2\kappa}) = (T^{-1} J_T^{*2})^{-\kappa} T^{-(\kappa-1)} T^{-1} \sum_{t=1}^T \mathbf{E}^*(|u_t^*|^{2\kappa}) |\tilde{\varepsilon}_t|^{2\kappa} = o_p(1),$$

for any $\kappa \in (1, 2]$, since $T^{-1} \sum_{t=1}^T |\tilde{\varepsilon}_t|^{2\kappa} = O_p(1)$ on account of $\mathbf{E}(u_t^4)$ being bounded uniformly in t . Thus, a standard subsequence argument and the classical functional central limit theorem for rowwise-independent triangular arrays allow us to deduce that

$$\begin{pmatrix} T^{-1/2} \sum_{t=T_1}^{T_2} u_t^* \\ T^{-3/2} \sum_{t=T_1}^{T_2} Y_{t-1}^* u_t^* \end{pmatrix} \rightsquigarrow_p \begin{pmatrix} \sigma \{W(r_2) - W(r_1)\} \\ \alpha \sigma \{r_2 W(r_2) - r_1 W(r_1) - \int_{r_1}^{r_2} W(s) ds\} \end{pmatrix}. \quad (\text{A.13})$$

Furthermore, putting $\xi_t^* := \sum_{j=1}^t u_j^*$, it is easy to verify that $T^{-3/2} \sum_{t=T_1}^{T_2} \xi_{t-1}^*$, $T^{-2} \sum_{t=T_1}^{T_2} \xi_{t-1}^{*2}$, and $T^{-5/2} \sum_{t=T_1}^{T_2} (t-1) \xi_{t-1}^*$ are of order $O_p^*(1)$. Hence, since $\tilde{\alpha} = \alpha + o_p(1)$, it follows that

$$T^{-2} \sum_{t=T_1}^{T_2} Y_{t-1}^* = \tilde{\alpha} T^{-2} \sum_{t=T_1}^{T_2} (t-1) + o_p^*(1) = \frac{1}{2} \alpha (r_2^2 - r_1^2) + o_p^*(1), \quad (\text{A.14})$$

and

$$T^{-3} \sum_{t=T_1}^{T_2} Y_{t-1}^{*2} = \tilde{\alpha}^2 T^{-3} \sum_{t=T_1}^{T_2} (t-1)^2 + o_p^*(1) = \frac{1}{3} \alpha^2 (r_2^3 - r_1^3) + o_p^*(1). \quad (\text{A.15})$$

Using (A.12), (A.13), (A.14), (A.15), and the conditional version of Slutsky's theorem, we may conclude, therefore, that

$$T^{3/2} \hat{\theta}^* \rightsquigarrow_p Z. \quad (\text{A.16})$$

Next, let $ADF^*(r_1, r_2) = (T^{3/2}\hat{\theta}^*)(T^{-3/2}\hat{\tau}^*)/\hat{\sigma}^*$ be the bootstrap version of $ADF(r_1, r_2)$,

where

$$\hat{\tau}^{*2} := \sum_{t=T_1}^{T_2} Y_{t-1}^{*2} - (T_2 - T_1 + 1)^{-1} \left(\sum_{t=T_1}^{T_2} Y_{t-1}^* \right)^2,$$

and

$$\hat{\sigma}^{*2} := (T_2 - T_1 - 1)^{-1} \sum_{t=T_1}^{T_2} \hat{u}_t^{*2}.$$

In view of (A.14) and (A.15),

$$T^{-3}\hat{\tau}^{*2} = \frac{1}{3}\alpha^2(r_2^3 - r_1^3) - \frac{1}{4}\alpha^2(r_2^2 - r_1^2)^2(r_2 - r_1)^{-1} + o_p^*(1) = \frac{1}{12}\alpha^2(r_2 - r_1)^3 + o_p^*(1). \quad (\text{A.17})$$

Moreover, it can be readily verified that $\hat{\sigma}^{*2} = (T_2 - T_1 - 1)^{-1} \sum_{t=T_1}^{T_2} u_t^{*2} + o_p^*(1)$, and so

$$\begin{aligned} \hat{\sigma}^{*2} - \sigma^2 &= (T_2 - T_1 - 1)^{-1} \sum_{t=T_1}^{T_2} (\nu_t^{*2} - 1)\tilde{\varepsilon}_t^2 + (T_2 - T_1 - 1)^{-1} \sum_{t=T_1}^{T_2} (\tilde{\varepsilon}_t^2 - \sigma^2) + o_p^*(1) \\ &= (T_2 - T_1 - 1)^{-1} \sum_{t=T_1}^{T_2} (\nu_t^{*2} - 1)\tilde{\varepsilon}_t^2 + o_p^*(1) \\ &= o_p^*(1). \end{aligned} \quad (\text{A.18})$$

The last equality is a consequence of $(T_2 - T_1 - 1)^{-1} \sum_{t=T_1}^{T_2} (\nu_t^{*2} - 1)\tilde{\varepsilon}_t^2 =: S^*$ satisfying

$$\begin{aligned} \mathbf{E}^*(S^{*2}) &= (T_2 - T_1 - 1)^{-2} \sum_{t=T_1}^{T_2} \tilde{\varepsilon}_t^4 \{\mathbf{E}^*(\nu_t^{*4}) - 1\} \\ &\leq (T_2 - T_1 - 1)^{-1} M_\nu \left((T_2 - T_1 - 1)^{-1} \sum_{t=T_1}^{T_2} \tilde{\varepsilon}_t^4 \right) = o_p(1), \end{aligned}$$

because of $\mathbf{E}^*[\tilde{\varepsilon}_t^2 \tilde{\varepsilon}_j^2 (\nu_t^{*2} - 1)(\nu_j^{*2} - 1)] = \tilde{\varepsilon}_t^4 \{\mathbf{E}^*(\nu_t^{*4}) - 1\} \delta_{tj}$ (δ_{tj} being Kronecker's delta) and $(T_2 - T_1 - 1)^{-1} \sum_{t=T_1}^{T_2} \tilde{\varepsilon}_t^4 = O_p(1)$, which implies, via the conditional version of Markov's inequality, that $S^* = o_p^*(1)$.

The results in (A.16), (A.17) and (A.18), together with the conditional version of Slutsky's theorem, ensure that

$$ADF^*(r_1, r_2) \rightsquigarrow_p \frac{\sqrt{3}}{(r_2 - r_1)^{3/2}} \left[(r_2 - r_1) \{W(r_2) + W(r_1)\} - 2 \int_{r_1}^{r_2} W(s) ds \right].$$

The proof is completed with the aid of the continuous mapping theorem by adapting the argument used in the proof of Theorem 1 of PSY to the bootstrap space carrying $\{Y_t^*\}$.

□

Under the conditions of Proposition A.2, the wild-bootstrap GSADF test is asymptotically correct, in the sense that $\mathbb{P}[GSADF(r_0) > G_\beta^*] \rightarrow \beta$ as $T \rightarrow \infty$, for any $\beta \in (0, 1)$, where G_β^* is the $(1 - \beta)$ -quantile of the conditional distribution of $GSADF^*(r_0)$, given \mathcal{Y} . This follows from Lemma 4.2 of Bücher and Kojadinovic (2019) upon noting that, on account of Propositions A.1 and A.2, and the absolute continuity of the distribution of Λ (cf. Lifshits (1983)), $|\mathbb{P}^*[GSADF^*(r_0) \leq x] - \mathbb{P}[GSADF(r_0) \leq x]| = o_p(1)$ uniformly in $x \in (-\infty, \infty)$.

Since the conditional quantile G_β^* is typically unavailable in practice, it may be approximated by the $\lfloor (B + 1)(1 - \beta) \rfloor$ -th largest of the values of B independent copies of $GSADF^*(r_0)$, say \hat{G}_β^* (cf. Section 4.3). By Lemma 4.2 of Bücher and Kojadinovic (2019), referring $GSADF(r_0)$ to \hat{G}_β^* yields an asymptotically correct test, in the sense that, for any $\beta \in (0, 1)$, $\mathbb{P}[GSADF(r_0) > \hat{G}_\beta^*] \rightarrow \beta$ as $T, B \rightarrow \infty$ under (A.5).

The asymptotic validity of the wild bootstrap in the case of the SADF test can be established in similar fashion. It is also straightforward to show that wild-bootstrap estimators of the sampling distributions of SADF and GSADF statistics are consistent when the DGP is given by (A.5) with $\alpha = 0$.

A.4 Simulation results

Tables A1–A7 show the proportion of 1000 artificial samples in which the null hypothesis of a unit root in $\ln P_t$ is rejected in favor of an explosive alternative (at the specified levels of significance) using the critical values in Table 1 of PSY. The DGP is given by (12) and (14), with $P_t = P_t^F$ for all t ; parameters other than the one indicated in each table are kept at their baseline values. Results for $\mu = 0.001$ and $\sigma_d = 0.01$ are reported in Table 3.

A.5 Ordinary-bootstrap simulation results

Table A8 shows the proportion of 1000 artificial samples in which the null hypothesis of a unit root in $\ln P_t$ is rejected in favor of an explosive alternative (at the specified levels of significance) using critical values obtained by non-calibrated and calibrated ordinary-bootstrap procedures ($B = 2N = 1000$). The DGP is given by (12) and (14), with the baseline parameter configuration and $P_t = P_t^F$ for all t .

T	SADF			GSADF		
	0.10	0.05	0.01	0.10	0.05	0.01
Asymptotic Critical Values						
100	0.0920	0.0510	0.0130	0.1140	0.0650	0.0260
200	0.1320	0.0700	0.0100	0.1340	0.0610	0.0180
400	0.1290	0.0560	0.0100	0.1190	0.0590	0.0130
800	0.2000	0.1030	0.0250	0.1710	0.0900	0.0310
Finite-Sample Critical Values						
100	0.1090	0.0580	0.0080	0.1220	0.0540	0.0190
200	0.1320	0.0720	0.0160	0.1130	0.0500	0.0090
400	0.1340	0.0560	0.0100	0.1360	0.0550	0.0110
800	0.1800	0.1020	0.0250	0.1320	0.0630	0.0190

Table A1: Rejection frequencies of recursive unit-root tests, $\mu = 0.0005$

T	SADF			GSADF		
	0.10	0.05	0.01	0.10	0.05	0.01
Asymptotic Critical Values						
100	0.2400	0.1540	0.0470	0.2630	0.1560	0.0680
200	0.3650	0.2180	0.0640	0.4070	0.2700	0.1110
400	0.421	0.283	0.102	0.484	0.362	0.125
800	0.5500	0.3920	0.1610	0.7710	0.6260	0.2940
Finite-Sample Critical Values						
100	0.2830	0.1800	0.0330	0.2720	0.1330	0.0380
200	0.3650	0.2230	0.0850	0.3650	0.2400	0.0640
400	0.429	0.283	0.108	0.520	0.358	0.100
800	0.5330	0.3790	0.1720	0.7010	0.5090	0.2130

Table A2: Rejection frequencies of recursive unit-root tests, $\mu = 0.002$

T	SADF			GSADF		
	0.10	0.05	0.01	0.10	0.05	0.01
	Asymptotic Critical Values					
100	0.3200	0.2240	0.0850	0.4060	0.3020	0.1250
200	0.4790	0.3400	0.0980	0.6160	0.4460	0.1820
400	0.5580	0.4070	0.1430	0.7560	0.5870	0.2580
800	0.6060	0.4500	0.1710	0.9330	0.8390	0.5140
	Finite-Sample Critical Values					
100	0.3760	0.2480	0.0670	0.4210	0.2450	0.0800
200	0.4790	0.3420	0.1400	0.5730	0.4050	0.1150
400	0.5600	0.4070	0.1510	0.7820	0.5810	0.2030
800	0.5860	0.4410	0.1840	0.8930	0.7450	0.3760

Table A3: Rejection frequencies of recursive unit-root tests, $\mu = 0.003$

T	SADF			GSADF		
	0.10	0.05	0.01	0.10	0.05	0.01
	Asymptotic Critical Values					
100	0.4080	0.2830	0.1160	0.5490	0.4250	0.1800
200	0.5400	0.3880	0.1360	0.7870	0.6330	0.3240
400	0.5800	0.3990	0.1490	0.8970	0.7650	0.3950
800	0.6600	0.5010	0.2440	0.9860	0.9400	0.7200
	Finite-Sample Critical Values					
100	0.4630	0.3200	0.0860	0.5590	0.3510	0.1180
200	0.5400	0.3880	0.1780	0.7440	0.5820	0.2100
400	0.5840	0.3990	0.1570	0.9160	0.7600	0.3200
800	0.6350	0.4920	0.2490	0.9610	0.8880	0.5720

Table A4: Rejection frequencies of recursive unit-root tests, $\mu = 0.004$

T	SADF			GSADF		
	0.10	0.05	0.01	0.10	0.05	0.01
	Asymptotic Critical Values					
100	0.2300	0.1410	0.0580	0.2620	0.1790	0.0570
200	0.3420	0.2240	0.0590	0.3850	0.2860	0.0960
400	0.4700	0.3000	0.0900	0.5150	0.3550	0.1140
800	0.5500	0.3860	0.1310	0.7430	0.5830	0.2840
	Finite-Sample Critical Values					
100	0.2820	0.1660	0.0380	0.2710	0.1380	0.0330
200	0.3420	0.2260	0.0730	0.3580	0.2510	0.0450
400	0.4750	0.3000	0.0950	0.5470	0.3530	0.0870
800	0.5280	0.3720	0.1370	0.6680	0.4790	0.1840

Table A5: Rejection frequencies of recursive unit-root tests, $\sigma = 0.005$

T	SADF			GSADF		
	0.10	0.05	0.01	0.10	0.05	0.01
	Asymptotic Critical Values					
100	0.0930	0.0570	0.0150	0.1120	0.0640	0.0200
200	0.1230	0.0760	0.0150	0.1350	0.0790	0.0240
400	0.1780	0.0870	0.0170	0.1540	0.0800	0.0120
800	0.2440	0.1300	0.0380	0.2300	0.1510	0.0420
	Finite-Sample Critical Values					
100	0.1210	0.0640	0.0110	0.1160	0.0510	0.0110
200	0.1230	0.0780	0.0250	0.1210	0.0640	0.0110
400	0.1810	0.0870	0.0170	0.1710	0.0780	0.0080
800	0.2280	0.1270	0.0400	0.1880	0.0960	0.0250

Table A6: Rejection frequencies of recursive unit-root tests, $\sigma = 0.015$

T	SADF			GSADF		
	0.10	0.05	0.01	0.10	0.05	0.01
Asymptotic Critical Values						
100	0.0910	0.0490	0.0180	0.1180	0.0760	0.0210
200	0.1270	0.0740	0.0070	0.1140	0.0570	0.0150
400	0.1330	0.0700	0.0130	0.1220	0.0590	0.0080
800	0.1880	0.0990	0.0300	0.1910	0.1040	0.0270
Finite-Sample Critical Values						
100	0.1170	0.0560	0.0150	0.1260	0.0600	0.0150
200	0.1270	0.0780	0.0110	0.0940	0.0480	0.0080
400	0.1370	0.0700	0.0140	0.1440	0.0570	0.0060
800	0.1710	0.0940	0.0310	0.1430	0.0740	0.0230

Table A7: Rejection frequencies of recursive unit-root tests, $\sigma = 0.02$

T	SADF			GSADF		
	0.10	0.05	0.01	0.10	0.05	0.01
Ordinary-Bootstrap Critical Values						
100	0.060	0.020	0.003	0.055	0.030	0.004
200	0.070	0.023	0.003	0.070	0.036	0.009
400	0.096	0.044	0.007	0.084	0.041	0.009
800	0.089	0.045	0.008	0.071	0.034	0.005
1600	0.108	0.058	0.008	0.102	0.050	0.009
Calibrated Ordinary-Bootstrap Critical Values						
100	0.088	0.032	0.004	0.085	0.041	0.005
200	0.092	0.044	0.008	0.091	0.050	0.011
400	0.104	0.054	0.009	0.103	0.049	0.010
800	0.093	0.053	0.008	0.088	0.042	0.004
1600	0.110	0.064	0.009	0.109	0.051	0.011

Table A8: Rejection frequencies of bootstrap-assisted recursive unit-root tests

References

- ALDOUS, D. (1978): “Stopping times and tightness,” *Annals of Probability*, 6, 335–340.
- ASTILL, S., D. I. HARVEY, S. J. LEYBOURNE, R. SOLLIS, AND A. M. R. TAYLOR (2018): “Real-time monitoring for explosive financial bubbles,” *Journal of Time Series Analysis*, 39, 863–891.
- BERAN, R. (1988): “Prepivoting test statistics: A bootstrap view of asymptotic refinements,” *Journal of the American Statistical Association*, 83, 687–697.
- BOURI, E., S. J. H. SHAHZAD, AND D. ROUBAUD (2019): “Co-explosivity in the cryptocurrency market,” *Finance Research Letters*, 29, 178–183.
- BÜCHER, A., AND I. KOJADINOVIC (2019): “A note on conditional versus joint unconditional weak convergence in bootstrap consistency results,” *Journal of Theoretical Probability*, 32, 1145–1165.
- CAMPBELL, J. Y., AND R. J. SHILLER (1989): “The dividend-price ratio and expectations of future dividends and discount factors,” *Review of Financial Studies*, 1, 195–228.
- CHEUNG, A., E. ROCA, AND J.-J. SU (2015): “Crypto-currency bubbles: An application of the Phillips-Shi-Yu (2013) methodology on Mt. Gox bitcoin prices,” *Applied Economics*, 47, 2348–2358.
- CORBET, S., B. LUCEY, AND L. YAROVAYA (2018): “Datestamping the Bitcoin and Ethereum bubbles,” *Finance Research Letters*, 26, 81–88.
- DRIFFILL, J., AND M. SOLA (1998): “Intrinsic bubbles and regime-switching,” *Journal of Monetary Economics*, 42, 357–373.
- ETIENNE, X. L., S. H. IRWIN, AND P. GARCIA (2014): “Bubbles in food commodity markets: Four decades of evidence,” *Journal of International Money and Finance*, 42, 129–155.
- EVANS, G. W. (1991): “Pitfalls in testing for explosive bubbles in asset prices,” *American Economic Review*, 81, 922–930.

- FANTAZZINI, D. (2016): “The oil price crash in 2014/15: Was there a (negative) financial bubble?,” *Energy Policy*, 96, 383–396.
- FROOT, K. A., AND M. OBSTFELD (1991): “Intrinsic bubbles: the case of stock prices,” *American Economic Review*, 81, 1189–1214.
- GREENAWAY-MCGREY, R., AND P. C. B. PHILLIPS (2016): “Hot property in New Zealand: Empirical evidence of housing bubbles in the metropolitan centres,” *New Zealand Economic Papers*, 50, 88–113.
- GÜRKAYNAK, R. S. (2008): “Econometric tests of asset price bubbles: Taking stock,” *Journal of Economic Surveys*, 22, 166–186.
- GUTIERREZ, L. (2013): “Speculative bubbles in agricultural commodity markets,” *European Review of Agricultural Economics*, 40, 217–238.
- HAFNER, C. M. (2020): “Testing for bubbles in cryptocurrencies with time-varying volatility,” *Journal of Financial Econometrics*, 18, 233–249.
- HARVEY, D. I., S. J. LEYBOURNE, R. SOLLIS, AND A. M. R. TAYLOR (2016): “Tests for explosive financial bubbles in the presence of non-stationary volatility,” *Journal of Empirical Finance*, 38, 548–574.
- HOMM, U., AND J. BREITUNG (2012): “Testing for speculative bubbles in stock markets: A comparison of alternative methods,” *Journal of Financial Econometrics*, 10, 198–231.
- HU, Y., AND L. OXLEY (2018): “Bubbles in US regional house prices: Evidence from house price–income ratios at the State level,” *Applied Economics*, 50, 3196–3229.
- KUAI, H., F. ALAJAJI, AND G. TAKAHARA (2000): “A lower bound on the probability of a finite union of events,” *Discrete Mathematics*, 215, 147–158.
- LIFSHITS, M. A. (1983): “On the absolute continuity of distributions of functionals of random processes,” *Theory of Probability and Its Applications*, 27, 600–607.
- LOH, W.-Y. (1987): “Calibrating confidence coefficients,” *Journal of the American Statistical Association*, 82, 155–162.

- LONG, W., D. LI, AND Q. LI (2016): “Testing explosive behavior in the gold market,” *Empirical Economics*, 51, 1151–1164.
- MONSCHANG, V., AND B. WILFLING (2021): “Sup-ADF-style bubble-detection methods under test,” *Empirical Economics*, 61, 145–172.
- PAVLIDIS, E., A. YUSUPOVA, I. PAYA, D. PEEL, E. MARTÍNEZ-GARCÍA, A. MACK, AND V. GROSSMAN (2016): “Episodes of exuberance in housing markets: In search of the smoking gun,” *The Journal of Real Estate Finance and Economics*, 53, 419–449.
- PETTENUZZO, D., R. SABBATUCCI, AND A. TIMMERMANN (2020): “Cash flow news and stock price dynamics,” *Journal of Finance*, 75, 2221–2270.
- PHILLIPS, P. C. B., AND S. SHI (2019): “Detecting financial collapse and ballooning sovereign risk,” *Oxford Bulletin of Economics and Statistics*, 81, 1336–1361.
- (2020): “Real time monitoring of asset markets: Bubbles and crises,” in *Handbook of Statistics: Financial, Macro and Micro Econometrics Using R*, ed. by H. D. Vinod, and C. R. Rao, vol. 42, pp. 61–80. North-Holland, Amsterdam.
- PHILLIPS, P. C. B., S. SHI, AND J. YU (2014): “Specification sensitivity in right-tailed unit root testing for explosive behaviour,” *Oxford Bulletin of Economics and Statistics*, 76, 315–333.
- (2015a): “Testing for multiple bubbles: Historical episodes of exuberance and collapse in the S&P 500,” *International Economic Review*, 56, 1043–1077.
- (2015b): “Testing for multiple bubbles: Limit theory of real-time detectors,” *International Economic Review*, 56, 1079–1133.
- PHILLIPS, P. C. B., Y. WU, AND J. YU (2011): “Explosive behavior in the 1990s Nasdaq: When did exuberance escalate asset values?,” *International Economic Review*, 52, 201–226.
- PHILLIPS, P. C. B., AND J. YU (2011): “Dating the timeline of financial bubbles during the subprime crisis,” *Quantitative Economics*, 2, 455–491.

WANG, X., AND J. YU (2022): “Bubble testing under polynomial trends,” *Econometrics Journal*, forthcoming.