



**RedNHE**

Red Nacional de  
Investigadores  
en Economía

# **Dynamic Arrangements in Economic Theory: Level-Agnostic Representations**

**Fernando Tohmé (UNS/CONICET)**

DOCUMENTO DE TRABAJO N° 274

Septiembre de 2023

Los documentos de trabajo de la RedNIE se difunden con el propósito de generar comentarios y debate, no habiendo estado sujetos a revisión de pares. Las opiniones expresadas en este trabajo son de los autores y no necesariamente representan las opiniones de la RedNIE o su Comisión Directiva.

The RedNIE working papers are disseminated for the purpose of generating comments and debate, and have not been subjected to peer review. The opinions expressed in this paper are exclusively those of the authors and do not necessarily represent the opinions of the RedNIE or its Board of Directors.

**Citar como:**

Tohmé, Fernando (2023). Dynamic Arrangements in Economic Theory: Level-Agnostic Representations. *Documento de trabajo RedNIE N°274*.

# Dynamic Arrangements in Economic Theory: Level-Agnostic Representations

Fernando Tohmé

Departamento de Economía - Universidad Nacional del Sur  
Instituto de Matemática de Bahía Blanca - CONICET

## Abstract

If Economics is understood as the study of the interactions among intentional agents, being rationality the main source of intentional behavior, the mathematical tools that it requires must be extended to capture systemic effects. Here we choose an alternative toolbox based on Category Theory. We examine potential *level-agnostic* formalisms, presenting three categories,  $\mathcal{PR}$ ,  $\mathcal{G}$  and an encompassing one,  $\mathcal{PR} - \mathcal{G}$ . The latter allows for representing dynamic rearrangements of the interactions among different agents.

## 1 Introduction

It is hard to define with precision the actual scope of Economics. Perhaps the best-known definition was given by Lionel Robbins (1932):

*Economics is the science which studies human behavior as a relationship between ends and scarce means which have alternative uses.*

While widely accepted, this characterization is unsatisfactory in many ways. In particular, for not taking into account crucial developments that reshaped the discipline in the last nine decades.

Accordingly, a more general definition could be

*Economics studies the interaction among intentional entities.*

This summarizes most if not all the research activities of contemporary economists. The term “entity”, which is introduced to refer to firms, institutions, and other non-human economic agents, covers the extension of economic analyses to all kinds of things able to exhibit agency, ranging from social groups to robots.<sup>1</sup>

---

<sup>1</sup>No wonder that other social scientists think that Economics is an “imperialist” discipline!

This extended notion of Economics, has been formalized in a rather narrower sense, using tools ranging from Calculus and Linear Algebra to Functional Analysis and Algebraic Topology. Modern Economic Theory as well as a good deal of Econometrics have been shaped using methods drawn from those fields. But the full meaning of the alternative characterization given above can only be captured by conceiving Economics as a system composed of other systems. While contemporary disciplines like Computer Science have fully embraced this view, economists have been reluctant to adopt it.

In this contribution we explore possible formalisms that may support the development of tools for an extended conception of Economics. While this is a wide-ranging project, we consider here two issues:

- How to deal with the complications inherent in attempts to sever different “local” interactions as if all the others remained fixed.
- How to scale up the solutions with the aggregation of the problems of interest.

Both issues reveal the need for a *level-agnostic* (or *continuous with respect to subagents*) Economic Theory. This paper lays the ground for a such model. We start by noting that Economics has a well-defined notion of agent defined in terms of a given preference relation over the space of alternatives. Then, the agent is said *rational* if she chooses the most preferred alternatives among those that are feasible for her.

In applications, it is customary to reduce the analysis to a subspace of the space of alternatives, simplifying the problem of making a decision. But this comes at the price of assuming the independence of the preferences over the subspace from the preferences over the rest of the larger space of alternatives.

In this initial version we first present a way of ensuring the consistency of the solutions found for the different subspaces. Then, another approach to the coordination of independent context is given, in this case involving games with shared players.

The final part of this paper presents a generalization, integrating both models, in which interactions are no longer fixed, but can evolve according to the inputs and outputs. In this as well as in the previous two models we apply the mathematical framework of Category Theory ([19]).

## 2 Mathematical Preliminaries

As is well-known, Category Theory has provided a framework without which most of the contemporary results in both Algebraic Geometry and Topology would not have been found [10]. As repeatedly shown in actual mathematical practice, the language of Set Theory remains insufficient for capturing perspicuously the nuances prevalent in those fields [14]. One reason is that unlike Set Theory the categorical approach allows for both the maximization of the “external” scope of its formal results *and* the controlled “internal” sensitivity to particular differences in content within the representation of mathematical structures. While Category Theory might thereby also seem to be a natural choice of a formal language for representing the decision-making problems outlined above, we have to note that Economics has been reluctant to adopt it.<sup>2</sup>

In this paper we draw heavily on the literature on Category Theory, although our results are clearly elementary. We will now present the basic concepts that will be used in subsequent sections. For further details and clarification, see the excellent presentations of Goldblatt ([9]), Barr & Wells ([3]), Adámek et al. ([2]), Lawvere and Shanel ([12]), Spivak ([18]), Fong and Spivak ([6]), Southwell ([17]) or Cheng ([4]).

A category  $\mathbf{C}$  consists of a set of *objects*,  $\text{Obj}$  and a class of *morphisms* between pairs of objects. Given two objects  $a, b \in \text{Obj}$  a morphism  $f$  between them is notated  $f : a \rightarrow b$ . Given another object  $c$  and a morphism  $g : b \rightarrow c$ , we have that  $f$  and  $g$  can be composed, yielding  $g \circ f : a \rightarrow c$  (COMPOSITION). Additionally, for every  $a \in \text{Obj}$ , there exists an *identity* morphism,  $\text{Id}_a : a \rightarrow a$ . Morphisms are required to obey two rules: (i) if  $f : a \rightarrow b$ ,  $f \circ \text{Id}_a = f$  and  $\text{Id}_b \circ f = f$  (IDENTITY); (ii) given  $f : a \rightarrow b$ ,  $g : b \rightarrow c$  and  $h : c \rightarrow d$ ,  $(h \circ g) \circ f = h \circ (g \circ f) : a \rightarrow d$  (ASSOCIATIVITY).

Examples of categories are **SET** (the objects are sets, and the morphisms are functions between sets), **TOP** (the objects are topological spaces and the morphisms continuous functions), **POrd** (the objects are preorders and the morphisms are order-preserving functions), **Vec** (the objects are vector spaces and the morphisms linear maps), etc.

The terseness of categories facilitates diagrammatic reasoning. A diagram in which nodes represent objects and arrows represent morphisms allows to establish properties of a category. Diagrams that *commute*, i.e. such that all different direct paths of morphisms with the same start and end nodes are identified (that is, compose to a common morphism), indicate relations similar to those that can be established by means of equa-

---

<sup>2</sup>Some notable exceptions are [8], [7], [1] and [16]. In turn, [5] presents arguments for the adoption of the categorical language in Economics.

tions.

Some of the most interesting constructions that can be defined in categories are *limits* and *colimits* (duals of limits). Any limit (or colimit) captures a *universal property* on a family of diagrams with the same basic shape. This basic shape is captured by a *cone*, that is, an object  $a$  and a family of arrows  $\{f_a^{b_j} : a \rightarrow b_j\}_{j \in \mathcal{J}}$ , such that for any pair  $j, l \in \mathcal{J}$ , if there exists a morphism  $\gamma_{jl} : b_j \rightarrow b_l$  we have that  $\gamma_{jl} \circ f_a^{b_j} = f_a^{b_l}$  (see Figure 1).




Figure 1: Commutative diagram

Then, given a class of cones of a given shape, a limit is an object  $L$  in this class such that for every other cone  $T$  in the class there exists a single morphism  $T \rightarrow L$  such that the resulting combined diagram commutes. For instance, consider a family of cones of the shape depicted in Figure 2.

$$a \longleftarrow \xrightarrow{f} X \xrightarrow{g} b$$

Figure 2: The limit of cones of this shape defines the product  $a \times b$

then, the limit is the *product*  $a \times b$  and with arrows  $p_1$  and  $p_2$ , the projections on the first ( $a$ ) and second ( $b$ ) components, respectively. For every other cone, with “apex”  $X$  there is a unique morphism  $! : X \rightarrow a \times b$  such that  $f = p_1 \circ !$  and  $g = p_2 \circ !$ .

Examples of colimits are *direct sums* (in SET, disjoint unions) and, somewhat confusingly called, *direct limits*, which in a self-contained description we will use to define *global solutions*.

Besides capturing interesting constructions common to many fields of Mathematics, Category Theory also provides tools for relating different categories to one another. This is achieved by means of mappings called *functors*. Given two categories  $\mathbf{C}$  and  $\mathbf{D}$  a functor  $F$  from  $\mathbf{C}$  to  $\mathbf{D}$  maps objects from  $\mathbf{C}$  into objects of  $\mathbf{D}$  as well as arrows from the former to the latter category such that, if

$$f : a \rightarrow b$$

in  $\mathbf{C}$ , then:

$$F(f) : F(a) \rightarrow F(b)$$

in  $\mathbf{D}$ . Furthermore  $F(g \circ f) = F(g) \circ F(f)$  and  $F(\text{Id}_a) = \text{Id}_{F(a)}$  for every object  $a$  in  $\mathbf{C}$ .

These functors are called *covariant*. Another class, that of *contravariant* functors, is such that, if

$$f : a \rightarrow b$$

in  $\mathbf{C}$ , then:

$$F(f) : F(b) \leftarrow F(a)$$

in  $\mathbf{D}$ . Of particular interest are the contravariant functors  $F : \mathbf{C} \rightarrow \mathbf{SET}$  (or a category of subsets of a given set), which are called *presheaves*. An intuitive interpretation is that given a morphism  $a \rightarrow b$  in  $\mathbf{C}$ , the morphism  $F(b) \rightarrow F(a)$  in  $\mathbf{SET}$  is the *restriction* of the “image” under  $F$  of  $b$  over the “image” of  $a$ . Given an object  $a$  in  $\mathbf{C}$ ,  $F(a)$  is called a *section* of  $F$  over  $a$ . This can be extended to any family  $B = \{b_j\}_{j \in \mathcal{J}}$  of objects in  $\mathbf{C}$ :  $F(B)$  is the section over  $B$ . In turn, given two families  $B \subseteq B'$  and the section over  $B'$ , namely  $F(B')$  we can find its restriction over  $B$ , denoted  $F(B')|_B$ , yielding  $F(B)$ .

Given a presheaf  $F : \mathbf{C} \rightarrow \mathbf{SET}$ , consider a class of objects  $B$  in  $\mathbf{C}$  and a cover  $\{K_j\}_{j \in \mathcal{J}}$  (i.e.  $B \subseteq \bigcup_{j \in \mathcal{J}} K_j$ ). Let  $\{k_j\}_{j \in \mathcal{J}}$  be a sequence such that  $k_j \in F(K_j)$  for each  $j \in \mathcal{J}$ . The presheaf  $F$  is said to be a *sheaf* if the following conditions are fulfilled:

- *Locality*: For every pair  $i, j \in \mathcal{J}$ ,  $k_i|_{K_i \cap K_j} = k_j|_{K_i \cap K_j}$  (i.e. the sections  $a_i, a_j$  coincide over  $V_i \cap V_j$ ),
- *Gluing*: There exists a unique  $\bar{b} \in F(B)$  such that  $\bar{b}|_{K_j} = k_j$  for each  $j \in \mathcal{J}$  (i.e. there exists a single object in the “image” of  $B$  that when restricted to each set in the covering yields the section corresponding to that set).

This brief review of Category Theory provides the basic concepts necessary for the analysis to be carried out in the rest of the paper.

Other notions will be introduced in the following sections.

### 3 Decision-making: Local vs. Global

The traditional characterization of decision-making under certainty by an individual is as follows. Let  $\mathcal{L}_i$  be a space of possible **options** that an agent  $i$  may select.<sup>3</sup> Each  $x \in \mathcal{L}_i$  is evaluated by means of a *utility* function,  $U_i : \mathcal{L}_i \rightarrow \mathfrak{R}$ . Given a family of constraints limiting the set of options open to the agent to  $\hat{\mathcal{L}}_i \subseteq \mathcal{L}_i$ , the goal of the agent is to find some  $\mathbf{x}^*$  that maximizes  $U_i$  over  $\hat{\mathcal{L}}_i$ . If we focus on the possible choices made by a single agent, we can drop the subindex  $i$  from the notation for  $\mathcal{L}$ ,  $\hat{\mathcal{L}}$  and  $U$ . We will reintroduce the dependence on the agents in the next sections to analyze the interaction between different agents.

In order to proceed, we first make some plausible assumptions. The space of options,  $\mathcal{L}$ , is presumed to be a (real) Hilbert space. That is, it is a complete metric space with an inner product. Furthermore, in order to ensure the existence of a  $\mathbf{x}^*$  we will also assume that  $\hat{\mathcal{L}}$  is a compact subset of  $\mathcal{L}$ , and that  $U$  is a continuous function. Within this very general framework, it is then straightforward to induce a category-theoretical representation of the global optimization of  $U$  over  $\hat{\mathcal{L}}$ , that is, of  $\mathbf{x}^*$  as a *direct limit*.

To begin, consider first a family  $\{L^k\}_{k=0}^{\kappa}$  of closed linear subspaces of  $\mathcal{L}$  and, for any given  $k$ , let us define the function

$$\text{Proj}_k : \mathcal{L} \rightarrow \bigcup_{k=0}^{\kappa} L^k$$

such that  $\text{Proj}_k(x) = x^k \in L^k$ , where  $x^k$  is the *projection* of  $x$  on  $L^k$ . The existence of such a projection is ensured by a straightforward application of the Linear Projection Theorem.<sup>4</sup>

The projector operator  $\text{Proj}_k$  will play a fundamental role in what follows. The intuition here is that we can think of each  $L^k$  as the options set of a local problem. Therefore, the projection of a global solution  $\mathbf{x}^*$  onto  $L^k$  will return the point in  $L^k$  which is the closest (i.e, the best!) to  $\mathbf{x}^*$ . Analogous approaches have been used successfully in several different contexts.<sup>5</sup>

---

<sup>3</sup>The meaning of these options depends on the context. If the agent is a consumer in a competitive market with a finite number of goods, she has to choose a vector of those commodities. In a planning problem, she has to select a plan specifying the amounts of resources used or consumed at each period of time.

<sup>4</sup>That is,  $|x - x^k| = \min_{y \in L^k} |x - y|$ , where  $|\cdot|$  is the norm of  $\mathcal{L}$ .

<sup>5</sup>See, among others, Luenberger [13], where these methods are employed to model pricing assets whose payoffs are outside the span of marketed assets



In case the projection does not return a local solution, we can still define an operator, which we call  $\Gamma_k : \hat{L} \rightarrow \hat{L}^k$  that formalizes the idea of a choice that is the closest to the projection (if it does not belong to the subspace):

$$\Gamma_k(x) = \{x^k \in \hat{\mathbf{X}}^k : x^k \in \operatorname{argmin}_{y \in \hat{\mathbf{X}}^k} |y - \operatorname{Proj}_k(x)|\}.$$

In some cases the global solution is not given, but must be sought by gluing together local ones “prospectively”, in the hope of producing (or better, abducting) a consistent global result. In order to formalize this broadly abductive method for seeking a global solution, we need to take a second, slightly deeper plunge into category theory and start with the definition of a category of local problems.

**Definition 1** Let  $\mathcal{PR}$  be the category of local problems, where

- $\operatorname{Obj}(\mathcal{PR})$  is the class of objects. Each one,  $s^k = \langle \hat{L}^k, u^k, \hat{\mathbf{X}}^k \rangle$  involves the maximization of the continuous utility function  $u^k$  over the compact set  $\hat{L}^k \subset L^k$ , a closed linear subspace of  $\mathcal{L}$ , yielding a family of solutions  $\hat{\mathbf{X}}^k$ .
- a morphism  $\rho_{kj} : s^k \rightarrow s^j$  is defined as  $\hat{L}^k \subseteq \hat{L}^j$ ,  $u^k = u^j|_{L^k}$  and  $\dim(L^k) \leq \dim(L^j)$ .<sup>6</sup> It follows from this definition that an identity morphism  $\rho_{kk} : s^k \rightarrow s^k$  trivially exists for every object  $s^k$ . Furthermore, given two morphisms  $\rho_{kj} : s^k \rightarrow s^j$  and  $\rho_{jl} : s^j \rightarrow s^l$  there exists their composition  $\rho_{jl} \circ \rho_{kj} = \rho_{kl}$ , since  $\hat{L}^k \subseteq \hat{L}^j \subseteq \hat{L}^l$ ,  $\dim(L^k) \leq \dim(L^j) \leq \dim(L^l)$  and by transitivity of the restrictions  $u^k = u^j|_{L^k}$  and  $u^j = u^l|_{L^j}$  we have that  $u^k = u^l|_{L^k}$ .




Figure 3: Morphism  $\rho_{kj}$  from  $s^k$  to  $s^j$ .

We can also define  $\mathcal{P}(\mathcal{L})$  as the category in which the objects are subsets of  $\mathcal{L}$  and a morphism between two objects  $f_{AB} : A \rightarrow B$  is defined as  $A \subseteq B$ .

Let us now define now a functor

$$\Sigma : \mathcal{PR} \rightarrow \mathcal{P}(\mathcal{L})$$

---

<sup>6</sup> $\dim(\cdot)$  yields the dimension of a subspace of  $\mathcal{L}$ .



Figure 4: Inclusion morphism representing  $A \subseteq B$ .

which assigns to a problem  $s^k = \langle \hat{L}^k, u^k, \hat{\mathbf{X}}^k \rangle$  the subset  $\Sigma(s^k)$  of  $\mathcal{L}$  defined by

$$\Sigma(s^k) = \{y \in \mathcal{L} \mid \Gamma_k(y) \in \hat{\mathbf{X}}^k\}$$

A section  $\sigma_k$  over  $s^k$  is simply the assignment of the elements of  $\Sigma(s^k)$  to  $s^k$ :

$$\sigma_k : s^k \mapsto \Sigma(s^k).$$

Given two problems,  $s^k = \langle \hat{L}^k, u^k, \hat{\mathbf{X}}^k \rangle$  and  $s^j = \langle \hat{L}^j, u^j, \hat{\mathbf{X}}^j \rangle$ , let us write  $s^k \triangleleft s^j$  iff there exists a morphism  $\rho$  in  $\mathcal{PR}$ ,  $\rho : s^k \rightarrow s^j$ . That is,  $s^k$  is a restriction of  $s^j$ .

Let us define  $r_k^j : \Sigma(s^j) \rightarrow \Sigma(s^k)$  such that to  $\Sigma(s^j)$  it assigns  $\Sigma(s^k)$ . Given a section over  $s^j$ ,  $r_k^j$  yields a section corresponding to its sub-problem  $s^k$ .

The following proposition then shows that the functor  $\Sigma$  possesses an important property that will be crucial for formalizing the possibility of patching up local problems and yielding a “larger” one:

**Proposition 1**  $\Sigma$  is a presheaf.

**Proof:**  $\Sigma : \mathcal{PR} \rightarrow \mathcal{P}(\mathcal{L})$  is a functor. We can analyze its behavior by means of  $r_k^j$ :

- For any  $s^k \in \text{Obj}(\mathcal{PR})$ , since  $s^k \triangleleft s^k$ ,  $r_k^k = \text{Id}_{\Sigma(s^k)}$ .
- If  $s^k \triangleleft s^j \triangleleft s^l$  then  $s^k \triangleleft s^l$ . Thus,  $r_k^j \circ r_j^l = r_k^l$ .

This means that  $\Sigma : \mathcal{PR} \rightarrow \mathcal{P}(\mathcal{L})$  is a contravariant functor. Or, in categorical terms, a presheaf.  $\square$

Consider now a family  $\{s^k = \langle \hat{L}^k, u^k, \hat{\mathbf{X}}^k \rangle\}_{k \in K} \subseteq \text{Obj}(\mathcal{PR})$ . It is said to be a *cover* of an object  $s^j = \langle \hat{L}^j, u^j, \hat{\mathbf{X}}^j \rangle$  of  $\text{Obj}(\mathcal{PR})$  if  $s^k \triangleleft s^j$  for each  $k \in K$  and  $\hat{L}^j \subseteq \bigcup_{k \in K} \hat{L}^k$ . That is, a problem  $s^j$  gets covered by the family  $\{s^k\}_{k \in K}$  if the domain of problem  $s^j$  is included in the union of the domains of the problems of the family and furthermore, each  $s^k$  is a

restriction of  $s^j$ .

The family of sections  $\{\sigma_k\}_{k \in K}$  is said to be *compatible* if for any pair  $k, l \in K$ , given  $\Sigma(s^k) = X^k$  and  $\Sigma(s^l) = X^l$ ,

$$\Gamma_k(X^k) \cap \Gamma_l(X^k) = \Gamma_k(X^l) \cap \Gamma_l(X^l)$$

Given a cover  $\{s^k\}_{k \in K}$  of a problem  $s^j$  with compatible sections,  $\Sigma$  is then a *K-sheaf* if there exists a unique  $\sigma_j = \Sigma(s^j)$  such that for each  $k \in K$ ,

$$\sigma_k = \sigma_j \cap \Gamma_k^{-1}(\hat{L}^k)$$

That is, intuitively,  $\Sigma$  is a K-sheaf if  $\sigma_j$  in fact “glues” together all the assignments  $\sigma_k$  in  $\mathcal{P}(\mathcal{L})$  within the more general framework of their compatibility. Finally, then, if  $\Sigma$  is a K-sheaf for every  $\{\sigma_k\}_{k \in K} \subseteq \text{Obj}(\mathcal{PR})$  it is called a *sheaf*.

**Example 1** Let  $\mathcal{L}$  to be  $\mathbb{R}^3$  (the three-dimensional real Euclidean space) and the utility function:

$$U(x, y, z) = 3 - 2x^2 - y^2 - 3z^2$$

to be maximized over  $\mathcal{L}$ . This yields a single global solution  $\hat{\mathbf{X}} = \{(0, 0, 0)\}$ .

Now consider two possible local problems:

- $L^1 = \{(x, y, z) : z = 0\}$ , with  $u^1(x, y, z) = U|_{L^1} = 3 - 2x^2 - y^2$  to be maximized over  $\hat{L}^1 = \{(x, y, 0) \in L^1 : x^2 + y^2 = 1\}$ , the unit circumference in  $L^1$ . The class of solutions for this problem is  $\hat{\mathbf{X}}^1 = \{(0, 1, 0), (0, -1, 0)\}$ .
- $L^2 = \{(x, y, z) : (x, y, z) \cdot (1, -1, 1) = 0\}$  (i.e. the linear subspace with normal vector  $(1, -1, 1)$ ), with  $u^2(x, y, z) = 3 - 3x^2 - 4z^2 - 2xz$ , the restriction of  $U$  on  $L^2$ , to be maximized over  $\hat{L}^2 = \{(x, y, z) : 2x^2 + 2z^2 + 2xz = 1\}$ , the intersection of the surface of the unit sphere in  $\mathbb{R}^3$  with  $L^2$ . Here the solution set is:  $\hat{\mathbf{X}}^2 = \{(-\sqrt{\frac{1}{3}}, -\frac{1}{2\sqrt{3}} - \frac{1}{2}, \frac{1}{2\sqrt{3}} - \frac{1}{2}), (\sqrt{\frac{1}{3}}, \frac{1}{2\sqrt{3}} + \frac{1}{2}, \frac{1}{2} - \frac{1}{2\sqrt{3}})\}$ .

It is easy to see that each solution of problem 1 minimizes the distance to the projection of the single global solution  $(0, 0, 0)$  on  $L^1$ . More precisely  $\Gamma_1(0, 0, 0) = \hat{\mathbf{X}}^1$ . The same is true for problem 2, since all points in  $L^2$  are at a Euclidean distance 1 from the global solution. So, in particular, the elements in  $\hat{\mathbf{X}}^2$  minimize the distance to the projection of  $(0, 0, 0)$  on  $L^2$  and thus,  $\Gamma_2(0, 0, 0) = \hat{\mathbf{X}}^2$ .

Given problems 1 and 2, denoted  $s^i = \langle \hat{L}^i, u^i, \hat{X}^i \rangle$  for  $i = 1, 2$ , we add a new problem  $s^0$ , which is the optimization of  $U$  over the surface of the three-dimensional sphere  $\hat{L}^0 = \{(x, y, z) : x^2 + y^2 + z^2 = 1\}$  and thus,  $\hat{X}^0 = \{(0, 1, 0), (0, -1, 0)\}$ . Suppose that these are the only objects in  $\mathcal{PR}$ . We define  $\Sigma : \mathcal{PR} \rightarrow \mathcal{P}(\mathcal{L})$ , summarized by the following table (each row being a section  $\sigma_i$ ,  $i = 0, 1, 2$ ):

Problems	$a_1$	$b_1$	$a_2$	$b_2$
$s^1$	X	–	X	–
$s^2$	–	X	–	X
$s^0$	X	–	X	–

The range of  $\Sigma$  is based only of four elements in  $\mathcal{L}$ :

$$a_1 = (0, 1, 0) \quad a_2 = (0, -1, 0)$$

and

$$b_1 = \left(-\sqrt{\frac{1}{3}}, -\frac{1}{2\sqrt{3}}, -\frac{1}{2}, \frac{1}{2\sqrt{3}}, -\frac{1}{2}\right) \quad b_2 = \left(\sqrt{\frac{1}{3}}, \frac{1}{2\sqrt{3}}, \frac{1}{2}, \frac{1}{2}, -\frac{1}{2\sqrt{3}}\right)$$

where  $a_1$  and  $a_2$  are the  $\mathbb{R}^3$  solutions of problems  $s^0$  and  $s^1$  while  $b_1$  and  $b_2$  are those of  $s^2$ .

It is easy to check that  $s^i \triangleleft s^0$  for  $i = 1, 2$ , since on one hand each problem  $s^i$  can be seen as the maximization of  $U$  restricted to subsets of the domain of problem  $s^0$ . On the other hand,  $\Sigma(s^0)$  restricted to each  $s^i$  yields  $\Sigma(s^i)$ . In fact, for  $s^1$  it is clear that this is the case. For  $s^2$ , let us note that  $b_1, b_2$  are the solutions of the problem  $s^0$  restricted to  $\hat{L}^2$ , seen as the inverse projection over the surface  $\hat{L}^0$ .

Furthermore,  $\{\sigma_1, \sigma_2\}$  is a compatible family of sections. Notice that  $\hat{L}^1 \cap \hat{L}^2$  does not include the solutions of either problem. But then the projections of  $\hat{X}^1$  and  $\hat{X}^2$  on  $\hat{L}^1 \cap \hat{L}^2$  are both  $\emptyset$ , and thus the sections satisfy, trivially, the compatibility condition. This means that  $\Sigma$  satisfies the sheaf condition.

Summarizing the discussion up to this point, we can say that given a category of problems  $\mathcal{PR}$  over a space  $\mathcal{L}$ , it is typically desirable to be able to obtain a sheaf  $\Sigma : \mathcal{PR} \rightarrow \mathcal{P}(\mathcal{L})$ , such that for any problem  $s^j$ , covered by any compatible family of sub-problems,  $\{s^k\}_{k \in K}$ ,  $\Sigma(s^j) \cap \Gamma_k^{-1}(\hat{L}^k) = \Sigma(s^k)$  for  $k \in K$ .

## 4 A Categorical Representation of Games

Let us now consider, instead of the coordination of different local decision problems, the coordination of games. That is, decision problems involving several agents, instead of a

single one. Thus, the approach discussed in this section generalized the sheaf-theoretical framework presented above.

Let us consider a category  $\mathcal{G}$  of *games*. Each object  $G$  in the category corresponds to a game  $G = \langle (I_G, S_G, \mathbf{O}_G, \rho_G), \pi_G \rangle$ , where

- $(I_G, S_G, \mathbf{O}_G, \rho_G)$  is a game form:
  - $I_G$  is the class of players.
  - $S_G = \prod_{i \in I_G} S_i^G$  is the *strategy set* of the game, where  $S_i^G \subseteq S_i$  is the set of strategies that player  $i$  can deploy in game  $G$ , for each  $i \in I_G$ .<sup>7</sup>
  - $\mathbf{O}_G$  is the class of *outcomes* of the game and  $\rho_G : S_G \rightarrow \mathbf{O}_G$  is a one-to-one function that associates each profile of strategies in the game with one of its outcomes.
- $\pi_G = \prod_{i \in I} \pi_i^G$  is a *profile of payoff functions*, where  $\pi_i^G : \mathbf{O}_G \rightarrow \mathbb{R}^+$  is the payoff function of player  $i$  in game  $G$ , for each  $i \in I_G$ .

A game is defined in terms of the interactions of *players*. Each player can be seen as described in terms of the strategies she can play and the payoffs she can receive from the results of her action (jointly with those of the other players).

We can define a category  $\mathcal{G}$ , where the objects are games. Given two games

$$G = \langle (I_G, S_G, \mathbf{O}_G, \rho_G), \pi_G \rangle \quad \text{and} \quad G' = \langle (I_{G'}, S_{G'}, \mathbf{O}_{G'}, \rho_{G'}), \pi_{G'} \rangle,$$

a morphism of games

$$G \rightarrow G'$$

is such that:

- $I_G \subseteq I_{G'}$ .
- $S_i^G \subseteq S_i^{G'}$  for each  $i \in I_G$ .
- There exist two functions, an inclusion  $p_{\mathbf{O}_G}^{\mathbf{O}_{G'}} : \mathbf{O}_{G'} \hookrightarrow \mathbf{O}_G$  for  $\mathbf{O}_{G'} \subseteq \mathbf{O}_G$  and a projection  $p_{S_G}^{S_{G'}} : S_{G'} \rightarrow S_G$ , i.e.  $p_{S_G}^{S_{G'}}(s_1^{G'}, \dots, s_i^{G'}, \dots, s_{|I_{G'}|}) \in \prod_{i \in I_G} S_i^{G'} = S_G$ . These functions verify the following condition:

$$\text{– For every } s' \in S_{G'}, s = p_{S_G}^{S_{G'}}(s') \in S_G \text{ is such that } \rho_G(s) = p_{\mathbf{O}_G}^{\mathbf{O}_{G'}}(\rho_{G'}(s')).$$

---

<sup>7</sup> $S_i$  is the set of all the strategies that player  $i$  can play in the games in which she participates.

Thus, if a morphism  $G \rightarrow G'$  exists,  $G$  can be conceived as a *subgame form* of  $G'$ .

To complete the characterization of  $\mathcal{G}$  notice that it is immediate that we can define *pushouts* and an *initial object* in this category:

- **Pushouts:** Consider three objects  $G, G'$  and  $G''$  and morphisms  $G \xrightarrow{f} G'$  and  $G \xrightarrow{g} G''$ . Then, take the coproduct of  $G'$  and  $G''$ , denoted  $G' + G''$ , obtained as the direct sums of the strategies sets and the outcomes of both games. By identifying the subgame forms of  $G'$  and  $G''$  corresponding to  $G$  we obtain the *pushout* of

$$G' \xleftarrow{f} G \xrightarrow{g} G''$$

- **Initial object:** Consider the *empty game*  $G^\emptyset$ , where  $I_{G^\emptyset} = \emptyset$  and consequently  $S_{G^\emptyset} = \emptyset$  and  $\mathbf{O}_G = \emptyset$  (thus  $\pi_{G^\emptyset}$  must be the empty function). It is immediate to see that  $G^\emptyset \rightarrow G$  for every  $G$  in  $\mathcal{G}$ .

Then we have

**Proposition 1**  $\mathcal{G}$  is a category with colimits.

Since  $\mathcal{G}$  is a category with colimits we can define *cospans* in it. Consider again three objects  $G, G'$  and  $G''$  and two morphisms  $G \xrightarrow{f} G'' \xleftarrow{g} G'$ . This is called a cospan from  $G$  to  $G'$ . The interpretation of such a cospan is that  $G$  and  $G'$  are subgame forms of the same game ( $G''$ ).

We can conceive each game  $G$  in  $\mathcal{G}$  as a *box*,  $G = (\text{in}^G, \text{out}^G)$ , where  $\text{in}^G$  and  $\text{out}^G$  are, respectively *input* and *output* ports.  $\text{in}^G$  has type  $\mathbf{O}_G$ , i.e. the input is an outcome of  $G$ . In turn, the  $\text{out}^G$  port has type  $S_G$ , being each output a profile in  $G$ .

Notice that each player  $i$  can be conceived as a game  $(\text{in}^i, \text{out}^i)$ , where  $\text{in}^i$  has type  $\cup_{G:i \in I_G} \mathbf{O}_G$  and  $\text{out}^i$  has type  $S_i$ .

Up to this point, our definition of morphisms in  $\mathcal{G}$  does not involve the payoffs. They can be incorporated by redefining the games as *modal boxes*, in which an additional component are the *internal states* of the game. More precisely, given any  $G$  and the class of its internal states,  $\Sigma_G$ , we can identify  $G$  as a triple  $\langle \text{in}^G, \text{out}^G, \Sigma_G \rangle$ , associated to two correspondences:

- **payoff:**  $\phi_G^1 : \bar{\text{in}}^G \times \Sigma_G \rightarrow \mathbb{R}^{+O_G}$ , such that for the vector  $o \in \bar{\text{in}}^G$  (the vector of all possible inputs of  $G$ , each entry being an outcome of the game) and state  $\sigma$ ,  $\phi_G^1(o, \sigma) = (\pi_G^i(o))_{o \in O_G}$ . That is, it yields the vector of payoffs corresponding to all the outcomes of  $G$ .
- **choice:**  $\phi_G^2 : \Sigma_G \rightarrow \bar{\text{out}}^G$ , such that for any state  $\sigma$ ,  $\phi_G^2(\sigma) = s \in \bar{\text{out}}^G$  (the class of all possible strategy profiles in  $S_G$ ) is a profile of strategies that may be chosen at that state.

Particularly relevant for our analysis is the definition of the internal states of each player  $i$ ,  $\Sigma_i$ . Consider a game  $G$  such that  $i \in I_G$ , and a sequence of morphisms in  $\mathcal{G}$

$$G_i^0 \rightarrow G_i^1 \rightarrow \dots \rightarrow G_i^{n-1} \rightarrow G_i^n$$

where  $G_i^0$  is a game in which  $i$  is the only player and  $G = G_i^n$ . We identify the state of player  $i$  when playing  $G$  as a sequence  $\sigma_G^i = \langle \sigma_0^i, \dots, \sigma_{n-1}^i \rangle$ , where  $\sigma_k^i \in \Sigma_{G_k^i}$ , for  $k = 0 \dots, n-1$ . Then, a distinguished object  $\sigma_*^i \in \Sigma_i$  is defined, such that  $\sigma_G^i$  is one of its initial segments.<sup>8</sup>

Therefore, for each game  $G$ ,  $\sigma_*^i$  can be instantiated yielding the corresponding state, and therefore the payoffs and the choices of player  $i$  in the game. The state  $\sigma_G$  of the entire game just obtains as the profile of states of its players.

A simple example is  $\sigma_{G^n}^i$  yielding as payoff for  $i$  the product of the payoffs she gets in the subgames of  $G^n$ . This case will be elaborated a bit more in Example 1, below.

We can define the category of cospans in  $\mathcal{G}$ , denoted  $\text{cospan}_{\mathcal{G}}$  which has a symmetric monoidal structure. Its objects are the same as those of  $\mathcal{G}$  and a morphism  $G \xrightarrow{h} G'$  is a cospan from  $G$  to  $G'$ , indicating that there exists a game of which  $G$  and  $G'$  are subgame forms. Thus, morphisms in  $\text{cospan}_{\mathcal{G}}$  are actually isomorphisms.

Given two morphisms in  $\text{cospan}_{\mathcal{G}}$ ,  $G \xrightarrow{f} G'$  and  $G' \xrightarrow{g} G''$  there exists a morphism  $G \xrightarrow{g \circ f} G''$  that obtains as a composition of the corresponding cospans.

The monoidal structure of  $\text{cospan}_{\mathcal{G}}$  is given by:

- The unit is  $G^\emptyset$ , the initial object in  $\mathcal{G}$ .
- The monoidal product of  $G$  and  $G'$ , is the coproduct  $G + G'$ .

---

<sup>8</sup>Thus,  $\sigma_*^i$  has a *forest* structure.

We now present a diagram language for open games. We start by considering the symmetric monoidal category  $\mathbf{W}_G$ . By definition, we have that:

$$\mathbf{W}_G = \text{cospan}_G$$

Each object, i.e. a game  $G$ , is seen as a  $\langle \text{in}^G, \text{out}^G, \Sigma_G \rangle$ -labeled *interface*, satisfying  $\phi_G^1$  and  $\phi_G^2$ . On the other hand, morphisms  $G \rightarrow C \leftarrow G'$ , are called  $\langle \text{in}, \text{out}, \Sigma \rangle$ -labeled *wiring diagrams*. The interpretation is that  $C$  is the overarching game that connects the subgames (not just the game forms)  $G$  and  $G'$ .

We write  $\psi : G_1, G_2, \dots, G_n \rightarrow \bar{G}$  to denote the wiring diagram  $\phi : G_1 + G_2 + \dots + G_n \rightarrow \bar{G}$ . We can, in turn see this as

$$G_1 + G_2 + \dots + G_n \xrightarrow{f} C \xleftarrow{\bar{f}} \bar{G}$$

which indicates that, being  $f$  and  $\bar{f}$  isomorphisms,

**Proposition 2**  $\bar{G}$  is the minimal game that includes the direct sum of  $G_1, \dots, G_n$  as a subgame.

## 5 Hypergraph Categories and Equilibria

We define a *hypergraph category*  $\langle \mathcal{G}, \text{Eq} \rangle$  with  $\text{Eq} : \mathbf{W}_G \rightarrow \prod_i S_i$ , such that, for every object  $G$  in  $\mathbf{W}_G$ ,  $\text{Eq}(G)$  is a class of vectors in  $\prod_{i \in I} S_i^G$ , the strategy set of game  $G$ . We assume that  $\text{Eq}(G)$  is a class of *equilibria* of  $G$ , for some notion of equilibrium (as for instance, dominant strategies equilibrium, admissible strategies, or Nash equilibrium).

**Example 2** Consider two games,  $G$  between players 1 and 2:<sup>9</sup>

	Bx	Bll
Bx	2, 1	0, 0
Bll	0, 0	1, 2

and  $G'$  between players 2 and 3:<sup>10</sup>


	C	D
C	2, 2	0, 3
D	3, 0	1, 1

<sup>9</sup>This is a *Battle of the Sexes* game, where  $S_1 = S_2 = \{\text{Bx}, \text{Bll}\}$ .

<sup>10</sup>A *Prisoner's Dilemma*, where  $S_2 = S_3 = \{C, D\}$ .



The corresponding wiring diagram is:



In red we have highlighted  $Eq(G) = \{(Bx, Bx), (Bll, Bll)\}$  and  $Eq(G') = \{(D, D)\}$ , where  $Eq$  corresponds to Nash equilibrium.<sup>11</sup>

Let us represent now  $G + G'$ . We start by building its corresponding game form. We obtain two tables, where the first one corresponds to player 3 choosing C:

	Bx-C	Bx-D	Bll-C	Bll-D
Bx	$o_{1,1}$	$o_{1,2}$	$o_{1,3}$	$o_{1,4}$
Bll	$o_{2,1}$	$o_{2,2}$	$o_{2,3}$	$o_{2,4}$

and another corresponding to player 3 choosing D:

	Bx-C	Bx-D	Bll-C	Bll-D
Bx	$o'_{1,1}$	$o'_{1,2}$	$o'_{1,3}$	$o'_{1,4}$
Bll	$o'_{2,1}$	$o'_{2,2}$	$o'_{2,3}$	$o'_{2,4}$

For instance,  $o_{11}$  indicates that 1 and 2 go to Box and 2 and 3 Cooperate. On the other hand,  $o'_{1,1}$  indicates that, again 1 and 2 go to Box, but while 2 keeps Cooperating, 3 Defects. The other entries can be interpreted likewise.

Suppose that the internal states of the players,  $\sigma_*^1, \sigma_*^2$  and  $\sigma_*^3$  are such that instantiated on  $G + G'$  yield the following payoffs and choices:

<sup>11</sup>Notice that here player 2, participates in two games.

If 3 chooses C:

	Bx-C	Bx-D	Bll-C	Bll-D
Bx	2,1 × 2,2	2,1 × 3,0	0,0 × 2,2	0,0 × 3,0
Bll	0,0 × 2,2	0,0 × 3,0	1,2 × 2,2	1,2 × 3,0

while if 3 chooses D:

	Bx-C	Bx-D	Bll-C	Bll-D
Bx	2,1 × 0,3	2,1 × 1,1	0,0 × 0,3	0,0 × 1,1
Bll	0,0 × 0,3	0,0 × 1,1	1,2 × 0,3	1,2 × 1,1

In words, players 1 and 3 keep the payoffs they get in the subgames, while 2 takes the product of the payoffs in  $G$  and  $G'$ . In red, we have highlighted the equilibria of  $G + G'$ , under this specification.

Let us define an operation  $\hat{\cup}$  such that given two equilibria  $s \in \text{Eq}(G)$  and  $s' \in \text{Eq}(G')$ , yields a new profile  $s - s' \in \text{Eq}(G) \hat{\cup} \text{Eq}(G')$  verifying that for each player  $i \in I_G \cap I_{G'}$ , a new strategy obtains combining  $s_i$  and  $s'_i$ , while in on all other cases the individual strategies are the same as in  $G$  and  $G'$ . Furthermore,  $\pi_i^{G \hat{\cup} G'}(s - s') = \pi_i^G(s) \times \pi_i^{G'}(s')$  for  $i \in I_G \cap I_{G'}$ .<sup>12</sup>

In our example, since  $\text{Eq}(G + G') = \{(Bx, Bx-D, D), (Bll, Bll-D, D)\}$ , we have that

$$\text{Eq}(G) \hat{\cup} \text{Eq}(G') = \text{Eq}(G + G').$$

This example illustrates the following claim:

**Proposition 3** For any pair of games  $G$  and  $G'$ ,  $\text{Eq}(G) \hat{\cup} \text{Eq}(G') = \text{Eq}(G + G')$ .

**Proof:** Trivial. If  $I_G \cap I_{G'} = \emptyset$ ,  $G + G' = G \cup G'$  with  $G \cap G' = \emptyset$ . Thus, each equilibrium of  $G + G'$  is just the disjoint combination of equilibria in  $G$  and  $G'$ .

If, on the other hand,  $I_G \cap I_{G'} \neq \emptyset$ , given  $i \in I_G \cap I_{G'}$ , her strategy set in  $G + G'$  is  $S_i^G \times S_i^{G'}$ , where  $S_i^G$  and  $S_i^{G'}$  are her strategy sets in  $G$  and  $G'$ , respectively. Now suppose that  $s_i^G$  and  $s_i^{G'}$

<sup>12</sup>An alternative yielding also Proposition 3 obtains if, instead, we take  $\pi_i^{G \hat{\cup} G'}(s - s') = \pi_i^G(s) + \pi_i^{G'}(s')$  for  $i \in I_G \cap I_{G'}$ .

are equilibrium strategies of  $i$  in the individual games but that  $(s_i^G, s_i^{G'})$  does not belong to an equilibrium in  $G + G'$ . Then, there exist an alternative combined strategy  $(\hat{s}_i^G, \hat{s}_i^{G'})$  such that on the new profile  $\pi_i$  yields a higher payoff, but since this equilibrium can be decomposed in two profiles, one in  $G$  and the other in  $G'$ , the payoff of  $i$  is the product of the payoffs over those two profiles. But then either  $\hat{s}_i^G$  yields a higher payoff than  $s_i^G$  or  $\hat{s}_i^{G'}$  yields a higher payoff than  $s_i^{G'}$  (recall that they are all positive real numbers). Thus, either  $s_i^G$  or  $s_i^{G'}$  is not an equilibrium in the corresponding game. Absurd.  $\square$

If we denote  $+$  the monoidal operation in  $\mathbf{W}_{\mathcal{G}}$ , if we take  $\otimes = \hat{\cup}$  as monoidal operation in  $\prod_i S_i$ , Proposition 3 indicates that there exist a trivial *natural isomorphism*

$$\text{Eq}(G) \otimes \text{Eq}(G') \rightarrow \text{Eq}(G + G')$$

Furthermore, taking the unit in  $\prod_i S_i$  to be the empty set, we have also that  $\emptyset = \text{Eq}(G^\emptyset)$ , where  $G^\emptyset$  is the initial object in  $\mathcal{G}$  and thus in  $\mathbf{W}_{\mathcal{G}}$ .

We have that

**Proposition 4** *Eq is a lax monoidal functor.*

Thus, the corresponding algebra allows to associate the composition of games with the equilibria of the components.

Proposition 4 depends critically on the possibility of defining  $\otimes$  in terms of a function  $\mathbf{f}$ , defined as follows. Given a player  $i \in I_G \cap I_{G'}$ , a combined strategy  $s_i - s'_i$  is such that for  $s = (s_i, s_{-i}) \in \text{Eq}(G)$  and  $s' = (s'_i, s'_{-i}) \in \text{Eq}(G')$ , satisfying  $\pi_i(s - s') = \mathbf{f}(\pi_i^G(s), \pi_i^{G'}(s'))$  and with  $s - s' \in \text{Eq}(G + G')$ . As we saw above if  $\mathbf{f}$  is the arithmetic product or sum, Eq will be indeed a lax monoidal functor.

But this restricts the compositionality of games to just trivial cases. We are interested in more general and non-obvious cases. In order to do that consider an alternative characterization of the hypergraph category  $\langle \mathcal{G}, \text{Eq} \rangle$ :

$$\text{Eq} : \mathbf{W}_{\mathcal{G}} \rightarrow \prod_i S_i \times \bigcup_{G \in \text{Obj}(\mathcal{G})} \Sigma_G$$

Furthermore, we need another definition of  $\otimes$ :

$$\otimes : \left( \prod_i S_i \times \bigcup_{G \in \text{Obj}(\mathcal{G})} \Sigma_G \right) \times \left( \prod_i S_i \times \bigcup_{G \in \text{Obj}(\mathcal{G})} \Sigma_G \right) \rightarrow \prod_i S_i \times \bigcup_{G \in \text{Obj}(\mathcal{G})} \Sigma_G$$

such that given two games  $G$  and  $G'$  with  $s \in \prod_{i \in I_G} S_i$  and  $\sigma_G$ , and  $s' \in \prod_{i \in I_{G'}} S_i$  and  $\sigma_{G'}$  we have:

$$(s, \sigma_G) \otimes (s', \sigma_{G'}) = (\bar{s}, \sigma_{G+G'}) \in \prod_{i \in I_{G+G'}} S_i \times \Sigma_{G+G'}$$

where  $\bar{s} \in S_{G+G'}$  is a Nash equilibrium if and only if  $s$  and  $s'$  are Nash equilibria of  $G$  and  $G'$  respectively.

$\otimes$  is well-defined. To see this, just recall that, by definition  $G + G'$  obtains in terms of the game forms of  $G$  and  $G'$  (the strategy sets and the outcomes), allowing different possible internal states and thus payoffs. The view of games as boxes presented in Section 4 indicates that there exist sequences of internal states of games, in parallel to sequences of morphisms between games, allowing to define  $\sigma_{G+G'}$ , and thus payoffs that make  $\bar{s}$  a Nash equilibrium if  $s$  and  $s'$  are also equilibria.

We can see that  $\prod_i S_i \times \bigcup_{G \in \mathbf{oxObj}(\mathcal{G})} \Sigma_G$  with  $\otimes$ , defined as above can be seen as a monoidal category, with morphisms defined in terms of those of  $\mathcal{G}$ , with  $(\emptyset, \emptyset)$  as its initial object. It allows to define Eq in such a way that by definition:

**Proposition 5** *Eq is a lax functor satisfying  $Eq(G + G') = Eq(G) \otimes Eq(G')$ .*

## 6 A more general model

$\langle \mathcal{G}, Eq \rangle$ , in any of the two versions of Eq seems too rigid to capture the dynamics of economic interactions. A more flexible structure is needed.

Let us start with the category of *polynomial functors*, **Poly**:

- Its objects have the following general form:

$$p = \sum_{i \in I} y^{p[i]}$$

where each term  $y^{p[i]}$  is a functor with domain  $p[i]$  into **Set**. Each  $i$  can be conceived as a *problem* while  $p[i]$  is a set of its *solutions*.

- Given  $p = \sum_{i \in I} y^{p[i]}$  and  $q = \sum_{j \in J} y^{q[j]}$  a morphism  $\phi : p \rightarrow q$  is  $\phi = (\phi_{\rightarrow}, \phi_{\leftarrow})$  such that

- $\phi^\rightarrow : I \rightarrow J$  and,
- $\phi^\leftarrow : q[\phi^\rightarrow(i)] \mapsto p[i]$ .

That is,  $\phi$  sends problems of  $I$  into problems of  $J$  and then the corresponding solutions in  $q$  back to the solutions in  $p$ .

We can conceive any  $p \in \text{Ob}(\mathbf{Poly})$  as an *interface* between inputs and outputs, being the inputs problems and the outputs their solutions. There are different ways of creating new interfaces up from other interfaces. We focus on the following construction:

- $[p, q] = \sum_{\phi: p \rightarrow q} y^{\sum_{i \in I} q[\phi^\rightarrow(i)]}$ , an *internal hom* in  $\mathbf{Poly}$ . It can be seen as a process that takes as inputs (*problems*) the morphisms from  $p$  to  $q$  and as outputs (*solutions*) all the possible solutions to the images of  $p$  in  $q$ .
- Given  $[p, q]$ , a  $[p, q] - \mathbf{Coalg}$  is a category in which each object is triple  $\langle s, \rho, \mu \rangle$ :
  - $s \in S$ , where  $S$  is a space of *states*, capturing the dynamics of the interface,
  - $\rho : s \mapsto (\phi, i, q[\phi^\rightarrow(i)])$ . That is, it assigns to the current state one of the solutions in  $[p, q]$ ,
  - $\mu$  updates the state in response to that pattern, i.e.  $\mu(\phi, i, q[\phi^\rightarrow(i)]) = s' \in S$ .

Consider now a category  $\mathbf{Org}$  defined as follows:

- $\text{Ob}(\mathbf{Org}) = \text{Ob}(\mathbf{Poly})$  and,
- $\text{Morph}(\mathbf{Org}) = [p, q] - \mathbf{Coalg}$ .

This means that two interfaces (connecting problems with their solutions)  $p$  and  $q$  are related by dynamic procedures of reconnection between them.

Our generalized model, covering *both*  $\mathcal{PR}$  and  $\langle \mathcal{G}, \text{Eq} \rangle$  is a category  $\mathcal{PR} - \mathcal{G}$  based on  $\mathbf{Org}$  such that, briefly:

- for each object  $a$  it corresponds  $p_a$  in  $\mathbf{Org}$ ,
- for objects  $a_1, \dots, a_n, b$  there corresponds a  $[p_{a_1} \otimes \dots \otimes p_{a_n}, p_b] - \mathbf{Coalg}$  of states  $S_{a_1, \dots, a_n, b}$ .<sup>13</sup>
- Each object  $a$  has an *identity* morphism.
- Pairs of morphisms compose.

---

<sup>13</sup>The operation  $p_a \otimes p_b$ , where  $p_a = \sum_{i \in I} y^{p_a[i]}$  and  $p_b = \sum_{j \in J} y^{p_b[j]}$ , is such that for each problem  $(i, j) \in I \times J$  yields the solutions to  $i$  and  $j$ ,  $p_a[i]$  and  $p_b[j]$ .

The last two requirements indicate, roughly, that morphisms inherit the identity and compositionality properties of **Org**

**Theorem 1** *Both  $Ob(\mathcal{G}) \subseteq Ob(\mathcal{PR} - \mathcal{G})$  and  $Ob(\mathcal{PR}) \subseteq Ob(\mathcal{PR} - \mathcal{G})$ .*

**Proof:** *Each problem in  $\mathcal{PR}$  can be interpreted as an interface between the problem itself and its optimal solutions. The same applies to any interactive decision-making setting in  $\mathcal{G}$ .*

*More precisely, a local problem  $s^k \in Ob(\mathcal{PR})$  and a game  $G \in Ob(\langle \mathcal{G}, Eq \rangle)$  can be represented by polynomial functor  $p_{s^k}$  or  $p_G$ , respectively. In the former case,  $p_{s^k}$  is an interface between the specification of the local problem  $(\hat{L}^k, u^k)$  and its solutions  $\hat{X}^k$ . In the case of a game,  $p_G$  is an interface between the game  $G$  and its equilibria  $Eq(G)$ .*

*Each state in the morphism between two interfaces  $p_{s^k}$  and  $p_{s^j}$  represents a particular  $r_j^k : \Sigma(s^k) \rightarrow \Sigma(s^j)$  that sends a section of solutions over  $s^k$  to a corresponding section over  $s^j$ , yielding a sheaf.*

*Analogously, each state in the morphism between two interfaces  $p_G$  and  $p_{G'}$  represents a particular wiring, connecting the games  $G$  and  $G'$ , such that the equilibrium obtains by tensoring those of the two games.  $\square$*

Notice that neither  $\mathcal{PR}$  nor  $\mathcal{G}$  are subcategories of  $\mathcal{PR} - \mathcal{G}$ . While their objects are also objects of the latter, morphisms among them are not morphisms in  $\mathcal{PR} - \mathcal{G}$ , which support dynamic rearrangements of the relations between its objects. Thus,  $\mathcal{PR} - \mathcal{G}$  incorporates all the representational advantages of  $\mathcal{PR}$  and  $\mathcal{G}$ , adding the possibility of capturing the dynamics of actual systems.

The following two examples exhibit the representational power of  $\mathcal{PR} - \mathcal{G}$ :

**Example 3** ([15]): *Consider a Principal-Agent problem defined by two functions:*

$$\Phi_{\rightarrow} : X \times Y \times \mathbb{R} \rightarrow \mathbb{R} \text{ and } \Pi : X \times Y \times \mathbb{R} \rightarrow \mathbb{R}$$

*where:*

- *$X$  is the compact set of types of the Agent.*
- *$Y$  is the compact set of possible decisions made by the Agent.*
- *$\Phi_{\rightarrow}$  is continuous, strictly decreasing in the third argument.*

- $\Phi_{\rightarrow}$  is full range in the third argument:  $\Phi_{\rightarrow}(x, y, \cdot)[\mathbb{R}] = \mathbb{R}$  for every  $(x, y) \in X \times Y$ .
- $\Pi$  is continuous and increasing in the third argument.
- $\Pi$  is full range in the third argument:  $\Pi(x, y, \cdot)[\mathbb{R}] = \mathbb{R}$  for every  $(x, y) \in X \times Y$ .

Given a type  $x$  of the Agent, her decision  $y$  and  $v$ , the money transfer to the Principal,  $\Phi_{\rightarrow}(x, y, v) = u_A$  is the utility of the Agent, while  $\Pi(x, y, v) = u_P$  is the utility of the Principal.

An inverse generating function is

$$\Phi^{\leftarrow} : Y \times X \times \mathbb{R} \rightarrow \mathbb{R}$$

such that given  $u_A = \Phi_{\rightarrow}(x, y, \Phi^{\leftarrow}(y, x, u_A))$  there exists  $v = \Phi^{\leftarrow}(y, x, \Phi_{\rightarrow}(x, y, v))$ .

Given  $\lambda \in \mathbb{M}$ , the class of Borel measures over  $X \times Y$  and  $\underline{u}$ , a reservation utility of the Agent, the Principal's problem amounts to choosing  $\langle \lambda, \bar{u}_A, \bar{v} \rangle$  as to maximize

$$\int_X \int_Y \Pi(x, y, \Phi^{\leftarrow}(y, x, \bar{u}_A)) d\lambda(x, y)$$

s.t.  $\bar{v} = \Phi^{\leftarrow}(y, x, \bar{u}_A)$  and  $\bar{u}_A \geq \underline{u}$ .

This setting can be naturally represented by defining two objects in  $\mathcal{PR} - \mathcal{G}$ ,  $A$ , and  $P$  (the Agent and the Principal, respectively). The corresponding polynomial functors are:

- $p_P$  takes as input  $\underline{u}$  and returns the optimal values  $\lambda^*$ ,  $u_A^*$  and  $\bar{v}^*$ . That is,  $p_P = \sum_{\underline{u} \in \mathbb{R}} \mathbf{y}^{p_P[\underline{u}]}$ , such that  $p_P[\underline{u}] = \langle \lambda^*, u_A^*, \bar{v}^* \rangle$ .
- $p_A$  takes as input  $\bar{v}$  and returns her decision  $y$  and the Principal's utility  $u_P$ . That is,  $p_A = \sum_{\bar{v} \in \mathbb{R}} \mathbf{y}^{p_A[\bar{v}]}$ , such that  $p_A[\bar{v}] = \langle y, u_P \rangle$ .

Then, the entire problem can be understood in terms of the identity morphism of  $p_A \otimes p_P$ , yielding the adjunction between  $\Phi^{\rightarrow}$  and  $\Phi^{\leftarrow}$ .

A promising area of research in which  $\mathcal{PR} - \mathcal{G}$  could be relevant for the design of mechanisms:

**Example 4** ([11] [7]): Mechanisms<sup>14</sup> can be conceived as game forms. That is, each mechanism  $M$  can be represented as  $M = (I_M, S_M, \mathbf{O}_M, \rho_M)$  (see Section 4).

Each  $i \in I_M$  can be given different incentives according the environment  $\mathbf{e} \in E$  in which she interacts with the others. Each  $\mathbf{e} \in E$  will have an associated profile of payoff functions that

---

<sup>14</sup>Institutions as well.

correspond to the outcomes in  $M$ ,  $\pi_M^{\mathbf{e}}$ .

The task of a mechanism designer  $D$  is to assign to a given environment a mechanism  $M \in \mathbb{M}$ , in order to ensure a target  $\mathbf{o}^*$ . Thus, in  $\mathcal{PR} - \mathcal{G}$ ,  $D$  has an associated  $p_D = \sum_{\mathbf{e} \in E} y^{p_D[\mathbf{e}]}$  where

$$p_D[\mathbf{e}] = \{ \langle M, \pi_M^{\mathbf{e}} \rangle : M \in \mathbb{M} \text{ such that } s_M^* \in \text{Eq}(\langle M, \pi_M^{\mathbf{e}} \rangle) \text{ and } \rho(s_M^*) = \mathbf{o}^* \in \mathbf{O}_M \}$$

Each game form  $M \in \mathbb{M}$  constitutes a local problem. The polynomial corresponding to these problems is  $p_M$ . In turn, given the choice of Nature (represented by a constant polynomial  $p_E = E$ ), the whole problem can be described by a  $[p_D \times p_E, p_M]$ -coalgebra, where:

$$[p_D \times p_E, p_M] = \sum_{\phi: p_D \times p_E \rightarrow p_M} y^{\sum_{\mathbf{e} \in E} p_M[\phi^{\rightarrow}(\mathbf{e})]}$$

and  $p_M[\phi^{\rightarrow}(\mathbf{e})] = \langle M, \pi_M^{\mathbf{e}} \rangle$ .

## 7 Conclusions

This paper discussed the question of representing economic phenomena in terms of interactions among intentional agents. We resorted to the language of Category Theory and, in particular, constructions like *sheaves*, *hypergraph categories*, and *polynomial functors*.

The category defined in terms of the latter,  $\mathcal{PR} - \mathcal{G}$ , has as objects the interfaces between problems and their solutions, while the interaction among them is captured by coalgebras based on the internal homs of the interfaces. That is, sets of states that determine the arrangement of connections among the problems and their solutions. Furthermore, the connections are rearranged in response to the outputs obtained previously.

We intend to explore further this formalism and use it to represent specific economic problems. While a first step involves showing that  $\mathcal{PR} - \mathcal{G}$  can reformulate known models, the real gist of this development is to capture new phenomena, establishing their relations to the former.

## References

- [1] S. Abramsky and V. Winschel: **Coalgebraic Analysis of Subgame-Perfect Equilibria in Infinite Games without Discounting**, *Mathematical Structures in Computer Science* **27** (2017), 751-761.



- [2] J. Adámek, H. Herrlich and G. Strecker: *Abstract and Concrete Categories. The Joy of Cats* (2004), <http://www.itl.cs.tu-bs.de/>.
- [3] M. Barr and Ch. Wells: *Category Theory. Lecture Notes for ESSLLI* (1999), <http://www.ling.ohio-state.edu/>.
- [4] E. Cheng: *The Joy of Abstraction*, Cambridge University Press, Cambridge (MA), 2022.
- [5] R. Crespo and F. Tohmé, F.: **The Future of Mathematics in Economics: a Philosophically Grounded Proposal**, *Foundations of Science* **22** (2017), 677-693.
- [6] B. Fong and D.I. Spivak: *An Invitation to Applied Category Theory: Seven Sketches in Compositionality*, Cambridge University Press, Cambridge (MA), 2019.
- [7] S. Frey, J. Hedges, J. Tan, P. Zahn: **Composing Games into Complex Institutions**, *PLoS ONE* **18** (2023), e0283361.
- [8] N. Ghani, J. Hedges, V. Winschel and P. Zahn.: **Compositional Game Theory**, *Proceedings of the 33rd Annual ACM/IEEE Symposium on Logic in Computer Science* (2018), 472-481.
- [9] R. Goldblatt: *Topoi. The Categorical Analysis of Logic*, North-Holland, Amsterdam, 1984.
- [10] A. Hatcher: *Algebraic Topology*, Cambridge University Press, Cambridge (UK), 2002.
- [11] L. Hurwicz: **Institutions as Families of Game Forms**, *Japanese Economic Review* **47** (1996), 113–132.
- [12] F. W. Lawvere and S. Schanuel: *Conceptual Mathematics: a First Introduction to Categories* (second edition), Cambridge University Press, Cambridge (UK), 2009.
- [13] D. Luenberger: **Projection Pricing**, *Journal of Optimization Theory and Applications* **109** (2001), 1-25.
- [14] J-P. Marquis: *From a Geometrical Point of View: a Study of the History and Philosophy of Category Theory*, Springer-Verlag, Berlin, 2009.
- [15] G. Nöldeke and L. Samuelson: **The Implementation Duality**, *Econometrica* **86** (2018), 1283-1324.
- [16] V. Rozen and G. Zhitomirski: **A Category Approach to Derived Preference Relations in Some Decision-Making Problems**, *Mathematical Social Sciences* **51** (2006), 257–273.

- [17] R. Southwell and N. Gupta: *Categories and Toposes: Visualized and Explained*, KDP Publishing, 2021.
- [18] D. I. Spivak: *Category Theory for the Sciences*, MIT Press, Cambridge (MA), 2014.
- [19] F. Tohmé and M. Auday: **Categorical Economic Theory**, in V. Beker (ed.) *Alternative Approaches to Economic Theory*, Routledge, London, 2019.
- [20] F. Tohmé, G. Caterina and R. Gangle: **Local and Global Optima in Decision-Making: a Sheaf-Theoretical Analysis of the Difference between Classical and Behavioral Approaches**, *International Journal of General Systems* **46** (2017), 879-897.